


L2 SEVIN®

ONLY FOR SALE TO, STORAGE AND USE BY SERVICE PERSONS  
OR FOR AGRICULTURAL USES.  
(Statement For Over 1 Gallon)



KILLS  
CHINCH-BUGS, JAPANESE BEETLES, BAGWORMS,  
EARWIGS, LAWN MOTHS (Sod Webworms),  
and many other insects as listed on  
FLOWERS • VEGETABLES • FRUITS  
LAWNS • SHRUBS • ORNAMENTALS

ACTIVE INGREDIENT:	BY WEIGHT
Carbaryl (1-Naphthyl Methylcarbamate).....	21.5%
INERT INGREDIENTS:.....	78.5%
TOTAL.....	100.0%

Contains 2 Pounds of Carbaryl per gallon.

® Trademark of Union Carbide Corp. for active ingredient  
(1-Naphthyl Methylcarbamate)

EPA Reg. No. 904-241

EPA Est. No. 72-PA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta  
que la etiqueta haya sido explicado ampliamente.

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by  
touching back of the throat with finger. Call a physician  
or poison control center immediately. Do not induce  
vomiting or give anything by mouth to an unconscious or  
convulsing person.

IF INHALED: Remove victim to fresh air. If not breathing, give  
artificial respiration, preferably mouth-to-mouth.

IF IN EYES: Immediately flush eyes with plenty of water. Call a physician.

IF ON SKIN: Remove contaminated clothing. Wash affected area with soap  
and water. Launder clothes before reusing.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase  
inhibitor. Atropine administered intravenously is  
antidotal. Do not use 2-PAM, opiates, or cholinesterase  
inhibiting drugs.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Pratt-Gabriel Division  
MILLER CHEMICAL & FERTILIZER CORPORATION  
Hanover, Pennsylvania 17331

NET CONTENTS:	8 OUNCES LIQUID	5 GALLONS LIQUID
	1 PINT LIQUID	30 GALLONS LIQUID
	1 QUART LIQUID	55 GALLONS LIQUID
	1 GALLON LIQUID	

**LAWNS:** For Ants, Chinch Bugs, Earwigs, Fall Armyworms, Fleas, Leafhoppers, Sod Webworms (Lawn Moths), and Millipedes, apply 8 fl. oz. in 15-20 gallons of water evenly over each 500 sq. ft. of established lawn area. For best results, mow the lawn and apply immediately after rain or watering. Do not water for at least 2 days after application. Repeat in 2-3 weeks if necessary. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

**ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES:** For use on Roses, Zinnias, Gladioli, Carnations, Lilacs, Arborvitae, Junipers, Hydrangeas, Azaleas, Elms, Chrysanthemums, Maples, Oaks, Dogwood, Birch and Pines (DO NOT USE ON BOSTON IVY OR VIRGINIA CREEPER) use 4 teaspoonsful per gallon of water or 1 pint in 25 gallons (1:200 dilution) to control Apple Aphid, Bagworm, Birchleaf Miner, Blister Beetle, Boxelder Bugs, Boxwood Leafminer, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetle, Gypsy Moth, Japanese Beetle, June Beetle, Lace Bug, Leafhoppers, Leafrollers, Mealybugs, Mimosa Webworms, Oakleaf Miners, Orange Tortrix, Periodical Cicada, Plant Bugs, Puss Caterpillars, Psyllids, Rose Slug, Scale Insects, Tent Caterpillars, Thorn Bugs, Thrips (exposed), Willowleaf Beetles. Apply when insects first appear. Repeat weekly or as necessary throughout the season.

**ADULT MOSQUITO CONTROL:** For Home Grounds, Parks and Recreational Areas, use 1:200 dilution in hydraulic low pressure sprayers (4 teaspoonsful per 1 gallon or 1 pint to 25 gallons of water). Apply in early morning or evening when air is calm and adults are active. Spray to thoroughly cover low shade tree foliage, shrubbery, flower beds, grass and shaded areas around buildings where adult mosquitoes congregate (DO NOT APPLY TO WATER). Repeat at 7-10 day intervals or as necessary (DO NOT SPRAY BOSTON IVY OR VIRGINIA CREEPER). May kill shrimp and crabs. Do not apply where these are important resources.

(Statement For Over 1 Gallon)

**RE-ENTRY AND ADDITIONAL APPLICATION RESTRICTIONS FOR  
AGRICULTURAL CROPS REQUIRING FARMWORKER HAND LABOR TASKS**

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing, until sprays have dried. Protective clothing means at least a hat or other suitable head covering, a long-sleeved shirt and long-legged trousers or a coverall type garment (all closely woven fabric covering the body), shoes and socks.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product. Instruct workers of the area or fields that must be vacated and not re-entered until the sprays have dried. Acquaint workers with the information appearing on this label under "Hazards to Humans and Domestic Animals" and "Statement of Practical Treatment". When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: 1) The name of this product; 2) Date of application; 3) The statement: "Do not enter without appropriate protective clothing until sprays have dried."; 4) The information given under "Hazards to Humans and Domestic Animals" and "Statement of Practical Treatment" that is appropriate in case of accidental exposure.

**DIRECTIONS FOR USE ON AGRICULTURAL CROPS REQUIRING FARMWORKER HAND LABOR TASKS**

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

**VEGETABLE GARDEN SPRAY:** Mix 1 fl. oz. in 3 gallons of water to 650 sq. ft. (2 qts. in sufficient water for 1 acre). Spray when insects or damage appear and repeat at 7-10 day intervals when necessary. No post-treatment time limitations for the following host plants when material is applied in accordance with the directions: BEANS (Green, Lima, and Snap), Japanese and Mexican Bean Beetle, Tarnished Plant Bug, TOMATOES, EGGPLANTS AND PEPPERS: Leafhopper, Lace Bug, Tomato Fruitworm, PUMPKINS, SQUASH, CUCUMBERS, MELONS: Flea Beetle, Melonworms, Cucumber Beetle, Squash Bug, SWEET CORN: Corn Earworm, Japanese Beetle and Leafhoppers: Mix 1 1/2 fl. oz. in 3 gallons of water to cover 650 sq. ft. (2 qts. in sufficient water for 1 acre). Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. For European Corn Borer, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge. Do not apply within 7 days of harvest if used for feed or forage. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

**LIMITED WARRANTY:** BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH THE DIRECTIONS HEREWITH.

treated areas until spray has dried.

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Do not use when value of bees as pollinators is more important than insect control. Warn beekeepers to locate hives beyond bee flight range until one week after application or take equally effective precautions.

PROTECT FROM TEMPERATURES BELOW 32° F. CLOSE CONTAINER IMMEDIATELY AFTER USING TO PREVENT EVAPORATION OR DRYING. SHAKE WELL BEFORE USING.

READ ENTIRE LABEL BEFORE USING -- OBSERVE ALL CAUTIONS -- USE ONLY AS DIRECTED.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### STORAGE & DISPOSAL (for sizes 1 Gallon and under)

Do not contaminate water, food or feed by storage or disposal. Store in tightly closed, original container in a locked area away from children and domestic animals. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

#### STORAGE & DISPOSAL (for sizes over 1 Gallon)

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in tightly closed, original container in a locked area away from children and domestic animals. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers may be also disposed of by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### MIXING DIRECTIONS

L2 Sevin® is an extremely fine dispersion of carbaryl in water. Stir thoroughly or shake before measuring. Pour recommended amount of concentrate into nearly full spray tank or premix material with water in another container and add to half full spray tank. Add balance of water with agitation. Agitate or stir well frequently while making application.

SPRAY THOROUGHLY TO COVER ALL PLANT SURFACES