



WASP & YELLOW JACKET SPRAY

JET STREAMS UP TO 12 FOOT RANGE
AIDS IN REACHING NESTS

Contains Resmethrin: KILLS Wasps • Yellow Jackets • Bees •
Hornets • Tent Caterpillars • Spiders



JAN 3 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 904-24/p

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION—May be harmful if swallowed or absorbed through the skin. Avoid inhalation of vapors. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Wash hands before eating or smoking. Do not contaminate food or feed. Avoid storage near food or feed products. Do not use on humans or household pets. Remove pets, birds, and cover fish aquariums before spraying.

In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with an acceptable detergent solution and rinse with potable water to remove all traces of contamination. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

PHYSICAL OR CHEMICAL HAZARDS

Extremely flammable. Contents under pressure. Keep away from fire, sparks, and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in cool area away from heat and open flame. Store in tightly closed, original container in a locked area away from children and domestic animals. For Disposal, replace cap, and securely wrap original container in several layers of newspaper and discard in trash. Do not incinerate or puncture.

DIRECTIONS

FOR USE in and around hotels, motels, homes, greenhouses and lathouses, food handling establishments, packaging and processing plants, seed houses, commercial, industrial, office, institutional and storage buildings, dairies, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in buses, automobiles, trucks, boxcars and ship cargo holds. Also for use in dog kennels and horse stables. In horse stables, not for application if horses will be used for food. Remove protective cap. Aim spray opening away from person. Push button to spray.

FOR household and industrial use, indoors and around premises. INDOORS: To kill wasps, hornets, yellow jackets, bees, application should be made late in the evening when insects are at rest. Spray liberally into hiding and breeding places, particularly under rafters, contacting as many insects as possible. Saturate areas in and around nest openings. Repeat as necessary. To kill ants, cockroaches, spiders, earwigs, grain mites, sowbugs, crickets, beetles, silverfish, centipedes and millipedes; direct spray to contact these insects in cracks and crevices in floors, baseboards, around pipes, behind cabinets and other harborages. Spray these infested areas until thoroughly wet. Repeat as necessary. To kill ants; spray directly on insects, and around window frames, doors, ant trails and hills. To kill spiders; spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders.

OUTDOORS Apply as a contact spray. Spray directly on insects. Aim spray opening away from person. Avoid spraying plants or turf except to spray infested areas. Ants: same as directed above for indoor use. Wasps, Hornets, Yellow Jackets, Bees: same as directed for indoor use. Aim spray at nest openings in trees, bushes, under eaves, chimneys, cracks and crevices to saturate in and around the nest openings. Contact as many insects as possible. Repeat as necessary. Earwigs, Sowbugs, Crickets, Beetles, Silverfish, Centipedes, Millipedes: same as a contact spray as for indoor use. Tent caterpillars, applications should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Aim spray directly at tents and totally saturate on all sides. Repeat as necessary.

LIMITED WARRANTY BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

Manufactured For:
Pratt-Gabriel Division
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331

105M7.5MP

Prod. No. 14814

Copyright © 1983 Miller Chemical & Fertilizer Corporation