

AGRISECT

BRAND

DIAZINON 5 % Granular

904-291

Jan 2, 1969

EN

ACTIVE INGREDIENTS

O,O-diethyl O-(2-Isopropyl - 4 - methyl -6-
pyrimidinyl) phosphorothioate 5%

INERT INGREDIENTS 95%
100%

CAUTION KEEP OUT OF THE REACH OF CHILDREN
HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOOD &
FEED PRODUCTS. AVOID INHALATION OF DUST. AVOID
CONTACT WITH SKIN, EYES AND CLOTHING. WASH THOROUGHLY
AFTER HANDLING. MAY BE ABSORBED THROUGH SKIN.

FISH & WILDLIFE IN TREATED AREAS MAY BE HARMED.
DO NOT CONTAMINATE FISH PONDS OR OTHER BODIES OF
WATER BY DIRECT APPLICATION, CLEANING OF EQUIPMENT,
OR DISPOSAL OF WASTES AND CONTAINERS. DISPOSE OF
EMPTY BAGS BY BURYING IN A SAFE PLACE.

Net 50 lbs.

BEST DOCUMENT AVAILABLE

AGRISECT
CHEMICAL CORPORATION

DIRECTIONS FOR USE

APPLICATIONS TO THE SOIL SHOULD BE WORKED IN THE TOP FEW
INCHES OF SOIL BY DISKING.

ONIONS: WIREWORMS-apply 40-80 lbs per acre as a preplant
broadcast treatment.

ONIONS: ONION MAGGOTS-apply 20 lbs per acre in the
furrow with the seed at the time of planting.

BROCCOLI, CABBAGE, CAULIFLOWER: ROOT MAGGOTS: apply
40-60 lbs per acre over entire seedbed and disk into
top 3-4 inches before seeding.

LAWNS: LAWN CHINCH BUG-apply 10 lbs per 5000 square
feet of lawn. Repeat application in 7-10 days.
Following application, brush lawn to assure that all
granules reach the soil surface.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING
OF THIS PRODUCT NOT IN STRICT ACCORDANCE WITH
DIRECTIONS GIVEN HEREWITH.

U.S.D.A. REG. NO. 8343-

ROBINNSVILLE, NEW JERSEY • NEW YORK, NEW YORK