

DIAZINON EMULSIFIABLE CONCENTRATE (contains 4% Diazinon per gallon)

ACTIVE INGREDIENTS:
O, O-diethyl-O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate*..... 47.5%
Aromatic petroleum derivative solvent.... 26.2%
INERT INGREDIENTS..... 26.3%
Total.....100.0%

*Diazinon

WARNING

KEEP OUT OF THE REACH OF CHILDREN

May be fatal if swallowed! May be absorbed through skin! Do not breathe spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed or foodstuffs. DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Apply only under supervision of trained personnel. Use suitable protective clothing and devices when handling concentrate or when spraying extensively. For special applications, this may be diluted with diesel oil or kerosene. A slight haze may develop, but utility of insecticide will not be affected.

DIRECTIONS FOR USE
ROACHES AND ANTS

Prepare a 1/2-percent emulsion by slowly adding 1-1/4 fluid ounces (2-1/2 tablespoons) of concentrate to 1 gallon of water with constant stirring. Apply as a strip treatment to cracks, crevices, and other roach harborages, and to ant runways. Use coarse, wet spray and apply until surface is wet to point of run-off (approximately 1 gallon per 1000 square feet). Reapply only when reinfestation is noted. Do not use for space spray or for residual treatments covering extensive indoor surfaces such as entire bulkheads and overheads. Preferably do not use in rooms occupied by children under 1 year of age.

FLY CONTROL (NOT INTENDED FOR LIVING QUARTERS)

"Spot" bait sprays. Apply a spray containing 1/2 pint of the concentrate and 1 pound of sugar in 2-1/2 gallons of water with a knapsack or similar type sprayer to areas frequented by flies, such as doorways, around windows, and other areas where flies congregate. Repeat applications every 2 to 3 weeks as fly population warrants.

Liquid bait for sprinkling can application. Use 2 fluid ounces of the concentrate and 1 pound of sugar (or 2 cups of syrup or molasses), in 5 gallons of water and sprinkle lightly over the floor in areas frequented

by flies. Burlap bags may be laid out in those areas frequented by flies and sprinkled with this mixture. Application every day or two will generally be required.

BROWN DOG TICKS AND FLEAS
IN BUILDINGS

Prepare a 1/2-percent emulsion as for roaches and ants above. Thoroughly spray infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets, and localized areas of floors and floor coverings. Applications should be repeated as infestations warrant and as reinfestations occur. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this formulation.

When used in dwellings, care should be taken to avoid deposits which could be frequently contacted by children. Care should also be taken to avoid spotting of wallpaper and fabrics. DO NOT CONTAMINATE FOOD, FOOD CONTAINERS, OR COOKING UTENSILS.

ACCEPTED
904-397
DEC 17 1968
UNDER THE FEDERAL INSECTICIDE AND RODENT CONTROL ACT FOR ECONOMIC PESTICIDE REGISTRATION SUBJECT TO ATTACHED COMMENTS.

DATE OF PACK:

LOT NO:

CONTRACT NO:

STOCK NO: FSN 6840-782-3925

NET CONTENTS: 1 GALLON

BEST DOCUMENT AVAILABLE

USDA REG. NO.

CONVERTED CENTRAL INSTITUTE
A