

VINELAND®

Net Weight (av.)
2.2 Pounds
(1000 gms.)

FORMALDEGEN-F®

U.S. Patent No. 2,993,832

ACCEPTED

2/21/68

FEDERAL INSECTICIDE
AND RODENTICIDE ACT
TOXIC PERSON REGISTER
REG. NO. 905-34

With the application of heat, Formaldegen-F will liberate Formaldehyde gas plus a compound which will fluoresce under ultra violet light.

Formaldegen-F is for use on hatching eggs either on the farm or at the hatchery. The inclusion of the compound which will fluoresce under ultra violet light is for the purpose of identifying at a later time the fact that such eggs have been fumigated with Formaldehyde gas.

Active Ingredients:
 Paraformaldehyde 91.00%
 Inert Ingredients 9.00%
 (Inert Ingredients Includes
 a Fluorescent Compound)



U.S.D.A. Reg. No. 905-34

Made in U.S.A.

RE-565210

VINELAND POULTRY LABORATORIES
 VINELAND, NEW JERSEY



POISON



If Formaldehyde Gas Is Inhaled

Remove patient to fresh air immediately. At first sign of difficulty in breathing, start artificial respiration; oxygen should be administered only by trained personnel. Continue artificial respiration until advised by physician to stop.

If FORMALDEGEN Is Taken Internally

Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Give milk or white of egg beaten with water. Call a physician immediately.

CAUTION: FORMALDEGEN, if used on exposed electric coils or in the presence of an open flame, is flammable. Care should be taken at all times to prevent this from happening.

WARNING: Causes irritation of skin, eyes, nose and throat. Do not breathe vapor. Avoid contact with skin, eyes or clothing. In case of contact flush with plenty of water; for eyes, get medical attention. Do Not Re-use Container. Bury empty container at least 18 inches deep in an isolated location away from water supplies.

U.S.D.A. Reg. No. 905-34

Made in U.S.A.

R1-863210

VINELAND®

FORMALDEGEN-F®

Net Weight (av.)
2.2 Pounds
(1000 gms.)

U.S. Patent No. 2,993,832

ACCEPTED

2/21/68

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 905-34

With the application of heat, Formaldegen-F will liberate Formaldehyde gas plus a compound which will fluoresce under ultra violet light.

Formaldegen-F is for use on hatching eggs either on the farm or at the hatchery. The inclusion of the compound which will fluoresce under ultra violet light is for the purpose of identifying at a later time the fact that such eggs have been fumigated with Formaldehyde gas.

Active Ingredients:
Paraformaldehyde 91.00%
Inert Ingredients 9.00%
(Inert Ingredients Includes
a Fluorescent Compound)

DANGER

Keep Out of Reach of Children



POISON



See Antidote and Additional Warnings On Left Panel

VINELAND POULTRY LABORATORIES
VINELAND, NEW JERSEY

VINELAND®

FORMALDEGEN-F®

Net Weight (av.)
2.2 Pounds
(1000 gms.)

U.S. Patent No. 2,993,832

With the application of heat, Formaldegen-F will liberate Formaldehyde gas plus a compound which will fluoresce under ultra violet light.

Formaldegen-F is for use on hatching eggs either on the farm or at the hatchery. The inclusion of the compound which will fluoresce under ultra violet light is for the purpose of identifying at a later time the fact that such eggs have been fumigated with Formaldehyde gas.

Active Ingredients:
Paraformaldehyde 91.00%
Inert Ingredients 9.00%
(Inert Ingredients Includes
a Fluorescent Compound)



POISON

See Antidote and Additional
Warnings On Left Panel

1. Remove all animal life from area to be fumigated.
2. Formaldehyde gas is most effective when humidity is 60% or higher and temperature is 60° F. or higher. Provide extra moisture and heat when necessary.
3. Calculate number of cu. ft. to be fumigated.
4. Use 20 grams of Formaldegen-F for each 100 cu. ft. of incubator, hatcher or egg fumigating cabinet space. When filled level, the enclosed special measuring spoon will hold 7 grams of Formaldegen-F. Liberate gas by application at 425° F. from electric hot plate having no exposed coils.
5. Fumigate hatching eggs for 20 minutes then exhaust cabinet or use Formaldegen Neutralizer.
6. Use 1 level measuring spoonful (7 grams) of Formaldegen-F for each run of eggs in the Chick Supplies automatic egg fumigation cabinet as described in the cabinet manufacturers' instructions.
7. After completion of fumigation, examination of the eggs fumigated with Formaldegen-F under ultra violet light will show white fluorescence whereas untreated eggs will have a purple color. Fluorescence may not be uniform over the shell of the egg but this is not important because deposition of the fluorescent dye depends upon settlement on the exposed portion of the shells during the fumigation process.

MEASURING SPOON INCLUDED IN CANISTER

ACCEPTED


2/1/68

FEDERAL INSECTICIDE
AND RODENTICIDE ACT
TOXIC POISON REGISTER-

905-34

DANGER

Out of Reach of Children



VINELAND POULTRY LABORATORIES
VINELAND, NEW JERSEY