




DDVP-CONCENTRATE

ACTIVE INGREDIENTS

2, 2-Dichlorovinyl Dimethyl Phosphate	46.5%
Related Compounds	3.5%
Inert Ingredients	50.0%
TOTAL	100.0%



DANGER

KEEP OUT OF THE REACH OF CHILDREN

Poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get in eyes or on skin. Do not breathe fumes. Rapidly absorbed through skin! In case of contact, wash immediately with soap and water. Do not contaminate feed and food-stuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Wear natural rubber gloves, waterproof protective clothing and goggles. Keep all unprotected persons out of operational area. While applying the insecticide, or working in treated areas during posted period, wear a mask or respirator of a type passed by the U. S. Bureau of Mines for protection against dimethyl dichloro vinyl phosphate.

DO NOT USE, SPILL, POUR OR STORE NEAR
HEAT OR OPEN FLAME

SEE ANTIDOTE
and other precautions on left panel.

DIRECTIONS

USE BY TRAINED PERSONNEL ONLY

This material is specifically formulated for timed & metered aerosol spraying equipment. When transferring from container to spraying equipment, the operator should wear a face mask passed by U. S. Bureau of Mines for protection against dimethyl dichloro vinyl phosphate, natural rubber gloves, waterproof protective clothing & goggles. The transfer area should be well ventilated with exhaust fan.

POSTING: Treated warehouse or factory area must be posted at each entrance for a period of 8 hours. No persons allowed in warehouse or factory area during treatment except for emergency and they must wear gas mask passed by U. S. Bureau of Mines for protection against dimethyl dichloro vinyl phosphate, natural rubber gloves, waterproof protective clothing & goggles. Take shower or bath after exit from building.

EQUIPMENT: Dispense between 5 p.m. and midnight only. One nozzle should be used per 25,000 cubic feet.

PRODUCT: Only for use in warehouse or factory area containing tobacco stored in hogsheads or cases. Not for use or storage in or around the home.

CIGARETTE BEETLE AND TOBACCO MOTH (exposed stages): Apply at a daily dosage rate of one gallon/5,262,000 cubic feet (.5 grams/1,000 cubic feet) For air conditioned warehouse or factory area use one gallon/2,631,000 cubic feet. (one gram/1,000 cubic feet)

NOTICE

THE SELLER GUARANTEES THE INGREDIENTS AS STATED.

Dispose of container by washing with detergent and water
thoroughly. Then sell to a cooperage company.

NET CONTENTS GALLONS

Manufactured by

EPA Reg. No. 908-42
EPA Est. 908-DC-1

LEFT PANEL

LEFT PANEL

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE is the emergency antidote for DDVP Insecticide poisoning. Consult your physician about obtaining an adequate supply of 1/100-grain atropine tablets for emergency use. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See WARNING SYMPTOMS below).

Keep patient prone and quiet.

Start artificial respiration immediately if patient is not breathing.

Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred.

2 PAM is also antidotal and may be administered in conjunction with atropine.

Morphine is contraindicated.

Clear chest by postural drainage.

Observe patient continuously for 48 hours.

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.

Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.