

~~QUICK~~ **QA** ~~ACTION~~

POISONED BARLEY

• FOR THE CONTROL OF •
California, Douglas, Fisher
GROUND SQUIRRELS



**POISON
DANGER**



KEEP OUT OF REACH OF CHILDREN

See antidote and other warnings on back panel.

Active ingredient:

Strychnine (expressed as alkaloid) 0.30%
Inert ingredients 99.70%

NET WEIGHT 1 LB.

ACCEPTED
FEB 28 1968
UNDER THE FEDERAL INSECTICIDE
HERBICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 221-54 SUBJECT
TO ATTACHED COMMENT.

USDA Reg. No. 909-11

DIRECTIONS

Scatter **Quick Action Poisoned Barley** on the ground near the holes and on the runways where squirrels find their food. About 1 tablespoonful should be scattered at each spot to cover 1 to 2 square feet.

For best results in hot weather place bait in burrows.

In farmyards or pastures or wherever there is livestock the poisoned grain must be placed inside the squirrel burrow in locations not accessible to children, pets and domestic animals. Do not scatter bait on bare ground.

NOTE: Strychnine readily kills various birds and rodents and, if taken in sufficient amount, can kill livestock.



WARNING: Convulsive poison. Poisonous if swallowed. Wash thoroughly after using. Do not contaminate feed and foodstuffs. This product is poisonous to fish and wildlife. Do not contaminate any body of water nor apply



when conditions favor drift or runoff from the crop area treated. Birds feeding in treated areas may be harmed.

POISON — ANTIDOTE

ANTIDOTE If less than ten minutes have passed since the poison was taken, give a tablespoonful of salt in a glass of warm water. Have victim lie down in a quiet, darkened room and keep him warm. Call a **PHYSICIAN IMMEDIATELY!**