


# DRAT

KILLS RATS AND MICE

# DRAT



# KILLS RATS AND MICE

CAUTION:

KEEP OUT OF REACH OF CHILDREN

COOKE LABORATORY PRODUCTS

# DRAT

KILLS RATS AND MICE

COOKE LABORATORY PRODUCTS PICO RIVERA CALIFORNIA

# DRAT KILLS RATS AND MICE

# DRAT KILLS RATS AND MICE

COOKE LABORATORY PRODUCTS PICO RIVERA CALIFORNIA

ACCEPTED

JAN 25 1968

UNDE  
FUN  
F

TO USE — NO MIXING REQUIRED

USDA REG. NO. 909-65

### DIRECTIONS FOR USE

Note: DRAT (contains Prolin\*) is hereinafter referred to as bait.

Place bait in locations where rats and mice feed, water or travel.

Bait placements along walls, in corners and in concealed locations where mice and rats can feed unobserved are desirable.

Bait should be placed in locations not accessible to children, pets or domestic animals or in tamper-proof bait boxes.

Contaminated or spoiled bait should be replaced promptly.

Baiting should be continued until all signs of feeding have stopped.

**RAT CONTROL:** Four ounces of bait in each bait location should be minimum amount placed. An uninterrupted supply of bait should be maintained for at least 15 days.

**MOUSE CONTROL:** Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended, and an uninterrupted supply of bait for a period of not less than 15 days should be assured.

### CAUTION

Protect children, pets and domestic animals from bait. If swallowed by humans call a physician immediately. This material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood, and, upon repeated feedings, causes fatal hemorrhage in rodents. It also contains an antibacterial agent, sulfaquinoxaline, which inhibits bacteria that produces vitamin K, an antidote for warfarin. It thereby increases effectiveness against hard-to-kill rats and mice and gives better and more thorough control.

\*Prolin (warfarin plus sulfaquinoxaline) is manufactured under license from the Wisconsin Alumni Research Foundation, U. S. Patent No. 2,427,578, other patent applications pending.


The antibacterial agent, sulfaquinoxaline, increases the effectiveness of this bait and assures more thorough control of rats. Especially recommended for rat infestations which are difficult to control.



LABORATORY PRODUCTS • PICO RIVERA, CALIFORNIA

# DRAAT

Contains PROLIN a better, more effective warfarin rodenticide



## KILLS RATS AND MICE

ACTIVE INGREDIENTS  
WARFARIN 10% ALUMINACE-ORNYLBENZYL 2-HYDROXYCOUMARIN  
N-2-QUINOLACETAMIDE SULFAQUINOXALINE  
INERT INGREDIENTS

### CAUTION:

See other cautions on back panel

KEEP OUT OF REACH OF CHILDREN

\*Trademark Wisconsin Alumni Research Foundation

### COOKE LABORATORY PRODUCTS

**COOKE LABORATORY PRODUCTS · PICO RIVERA, CALIFORNIA**  
**RAT KILLS RATS**  
**AND MICE**

**DRAAT KILLS RATS**  
**AND MICE**