ACCEPTED

TF85 1970

FANGECINE OF PUDENTS OF FUT FANGECINE OF PUDENTS OF FUT FANGECINES NO. 100 PUDENTS OF A STORY

DIRECTIONS (Continued)

Do not spray fruits, edible foliage or food crops. Spray along fences, walls and curbs: In places that provide shelter and shade, loose boards, stones, cracks and crevices. Wet surface thoroughly. Repeat as required to control.

Ants, on ground: 1 measuring tablespoonful to 1 gallon. Spray freely on infested areas, along curbs, fences and walls, in damp shady places, cracks and crevices. Repeat as required to control.

WARNING: Keep out of reach of children and pets. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, wash immediately with soap and water. For eyes, flush with plenty of water and get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Do not use on household pets or humans. Avoid contamination of feed and foodstuffs. Do not apply to food crops.

Highly toxic to fish and wildlife. Keep out of any body of water. Birds feeding in treated areas may be harmed.

Keep children and pets off treated areas until this material has been washed into the soil and area is thoroughly dry.

DANGER - FLAMMABLE! Keep away from heat or open flame. Keep in a cool place, out of direct sunlight. When container is empty, wash thoroughly and destroy. Do not re-use the container for any purpose.



WARNING: KEEP OUT OF REACH OF CHILDREN See other warnings on back panel.

USDA REG. NO. 909-67

NET CONTENTS ONE QUART - TREATS 3200 SQUARE FEET



- 1. By contact
- 2. By vapor
- 3. By ingestion

Active ingredients: Ethylene dichloride Cyclonexanone Gamma isomer of benzene hexachloride from lindane Petroleum hydrocarbon solvent Tokaphene Inert ingredients **Chlorinated camphene (chlorine content 67% to 69%).

DIRECTIONS

DILUTE WITH WATER AS DIRECTED BELOW

Cutworms, Army Worms, Lucerne Moth, Fiery Skipper and Lawn Moth (Sod Webworm) Larvae, White Grubs, Millipedes, Fleas in Dichondra and Grass Lawns: 2 measuring tablespoonfuls (1 ounce) to gallon. Water lawn before treatment. Apply spray solution, 1 gal. to 100 sq. feet. evenly over lawn. Then spray with plain water to wash insecticide off grass into soil. Keep children or pets off treated lawn until it has completely dried. Allow 2 days to elapse before re-watering. Repeat if and when re-infestation occurs

Aphids, Climbing Cutworms, Plant Bugs (Lygus), Flea Beetles, Elm Leaf Beetles, Leafhoppers, Tent Caterpillars, Spiny Elm Caterpillars, and certain other leaf-eating caterpillars on ornamental trees, shrubs and bushes. I measuring tablespoonful to gallon. Spray at first signs of damage or appearance of pests. Thoroughly wet pests and all parts of plant. Repeat as required to control.

Grasshoppers, Earwigs, Crickets, Sowbugs: 1 measuring tablespoonful to gallon. Spray freely on sail surface and foliage of flowers, ornamentals, shade trees, and weeds in infested area.

•

DIRECTIONS Consinues

Ants, on process in the tenness and second state of the second se

WARNING: First Warning of the send of the

PANGED HAMMABIT -



WARNING: REPORT OF REACH OF CHILDREN

See other warnings on back panel

NI ! CONTENTS ONE QUART TREATS 3200 SQUARE FEET



: ·

- 1 By contact
- 2 By vapor
- 3 By ingustion

DIRECTIONS

the growing the state of the second

Lawn Moth of A A Lawns White Groups Monday Fras Dichondra of A Salar Sal

Aphids, Climbing Cutworms. Plant Bugs (Lygus), Fina Beetins, Em Leat Beetins, Leath ppers, Tent Caterphiars, Spiny Em Caterphilars, and the desired of the control of the c

COOKE CABORATORY PRODUCTS . PICD RIVERA . CALIFORNIA