

Food Crops and Fruit Trees: Consult your local farm advisor for dosages and timing recommendations. To avoid excessive residues at harvest do not exceed the recommended dosage rates or the time limitation between the last application and harvest.

When spraying Oranges, Lemons, Grapefruit, Limes, Apples, Pears, Grapes, *Figs, *Cherries, *Plums, *Prunes, Tomatoes, Melons, Cucumbers, Squash, Strawberries, Boysenberries, Raspberries, Loganberries, Blackberries do not apply later than 7 days before harvest.

*Do not repeat application within 30 days.

When spraying *Apricots, *Peaches, *Nectarines, Walnuts, Pecans, Filberts, Almonds (Do not feed treated hulls to livestock) do not apply later than 14 days before harvest.

*Do not repeat application within 30 days.

NOTE: Do not add or mix with other spray materials, such as lime mixtures or wettable powders.

Caution: Avoid repeated skin contact. Avoid inhalation of mist. Do not take internally. Avoid contamination of foodstuffs. Wash thoroughly after handling. Do not smoke while spraying. **Inflammable**—Keep away from flame. This product is poisonous to fish.

* © Rohm & Haas—Philadelphia



CAUTION: See other cautions on back panel.

KEEP OUT OF REACH OF CHILDREN

Active ingredient:
 1,1,-Bis (p-chlorophenyl) - 18.5%
 2,2,2-trichloroethanol 81.5%
 Inert ingredients

USDA Reg. No. 909-69

DIRECTIONS

Cooke Red Spider and Mite Spray — a specific miticide with a broad range of effectiveness against important horticultural and agricultural mites such as Red Spider, Two-spotted, Clover (Brown Almond), Tomato Russet, Citrus Red, Citrus Flat, Plum Nursery, Peach Silver, Desert, Tropical, Apple Rust, Cyclamen, Privet, and Six-spotted mites.

all mites

For spray solution add —

- 1 to 2 teaspoonfuls to 1 gallon water
- 1 to 2 tablespoonfuls to 3 gallons water
- 2 to 4 ounces to 12 gallons water
- 1 to 2 pints to 100 gallons water

Stir or agitate to mix.

Spray thoroughly to give full coverage of plant, especially the undersides of leaves. On sensitive or tender foliage use the lesser dosage.

Ornamentals, Flowers, Nursery Stock and Shade Trees: Apply when mites are present and repeat as necessary.

Clover Mites in Buildings: Use 2 tablespoonfuls to 3 gallons water. Spray walls, foundation and window-sills and plants and lawn at base of infested buildings. Repeat as required to control.

Indoor
Indoor

COOKE LABORATORY PRODUCTS • PICO RIVERA • CALIFORNIA

