

COOKE
SLUG-N-SNAIL GRANULES

May Be Used Around Fruits and Vegetables

Granules Kill Snails, Slugs, Earwigs, Sowbugs, Cutworms, Ants

ACTIVE INGREDIENTS:

Metaldehyde 3.00%
Carbaryl (1-naphthyl N-methylcarbamate) 5.00%
INERT INGREDIENTS 92.00%

Keep out of reach of children

WARNING

This pesticide may be fatal to children, dogs or other pets if eaten. Keep children and pets out of the treated area. Follow directions and use with care. See other warnings on back panel.

NET WT. 1 LB.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Keep out of reach of children. May be fatal if swallowed. Avoid contact with skin, eyes, or clothing. Do not allow the granules to contact edible portion of any food or feed crop. Wash thoroughly after using. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes: Flush eyes with water. Consult a physician if irritation persists. If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. If On Skin: Wash thoroughly with soap and water. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

This pesticide may be fatal to children and dogs or other pets if eaten. Keep out of reach of children and pets, poultry and livestock. Protect dogs from treated areas, since they may be attracted to this product when applied.

ENVIRONMENTAL HAZARDS


This pesticide is toxic to birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not contaminate water by cleaning equipment or disposal of wastes.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label. Heed directions and warnings. Water infested area thoroughly. Scatter SLUG-N-SNAIL GRANULES on ground as to cover a large area - do not place in piles or heaps. Apply at anytime of day, although it is preferable to treat in late afternoon. After application do not water treated area for 48 hours. For best results, apply in the evening, especially following rain or watering.

To kill slugs and snails in ornamental gardens, planted borders, beds



and ground covers, such as ivy or dichondra or grass lawns: Scatter —
SLUG-N-SNAIL GRANULES lightly on surface of infested area at rate
of 1 pound to 1,500 square feet. Apply under and around flowers and
shrubs, along edges of flower beds adjacent to border or ground cover
plants or lawn. In crevices, as between lawn and walk or curb, and on
ground at base of fences and walls in shaded locations.

VEGETABLES: including beans, tomatoes, peas, corn, sweet corn, carrots,
cucumbers, peppers, cabbage, squash, broccoli, lettuce,
radishes, artichokes, beets, brussel sprouts, spinach, cauliflower,
swiss chard, okra, rhubarb, celery, egg plant, head lettuce.

FRUITS: including strawberries, cane berries, blackberries, avocados,
cantaloupes, cherries, citrus, grapes, peaches, apples, and plums,
melons. Do not allow the granules to contact edible portion of any food
or feed crop. Apply 3 or 4 weeks during the growing season, or as
needed. DO NOT APPLY DIRECTLY ONTO PLANTS.

For hard-to-reach places or under dense foliage in shaded areas where
these pests may hide, place SLUG-N-SNAIL GRANULES in a cup or a similar
receptacle and broadcast, with a sweeping motion, over the soil surface.
Repeat in 10 days or when reinfestation occurs.

LAWN AND GENERAL GARDEN USE: for control of Ants, Cutworms, Earwigs,
Fall Armyworms, Millipedes, and Sowbugs: Apply as directed above using
1 pound of SLUG-N-SNAIL GRANULES to 250 square feet. Repeat as required
to control. For earwig control: Make applications particularly to
daytime hiding places, including lawns, around base of trees, shrubs,
along walks, fences and foundation of buildings. Keep children and pets
off treated areas until this material has been washed into the soil and
area is thoroughly dry. For best results, apply in the evening,
especially following rain or watering.

STORAGE AND DISPOSAL

Avoid contamination of feed, food-stuffs or water
supplies. Store in a dry cool place preferably in a locked storage
area. Do not reuse container. Discard container in trash.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: Buyer assumes all risks of use, storage or handling of this
material not in strict accordance with directions given herewith.

COOKE LABORATORY PRODUCTS
PORTLAND, OREGON 97218
EPA Reg. No. 909-83
EPA Est. 802-OR-1
79132

