

COOP[®]

WITH 1% VAPONA

DIRECTIONS

Dairy and Beef Cattle — To control face flies, horn flies, stable flies and mosquitoes, apply 1 to 2 fluid oz. per animal, as a mist spray daily. Spray to thoroughly cover all surfaces of the animal including the legs, but not to wet the skin. Spray the animal's head (cheeks, muzzle and forehead) each morning as protection against face flies. Do not use over 2 fluid oz. per animal per day.

Fogging: Dairy Cattle, Beef Cattle and Barns — To control horn flies and stable flies. Use a barn fogger to apply 2 fl. oz. per 1,000 cu. ft. Before fogging, reduce the air movement by closing doors and windows. Open and air out the barn about 5 minutes after treatment. Do not contaminate feed, feedstuffs, milk or milk utensils. Do not fog animals that have received a direct spray application within 8 hours.

Poultry Houses — Apply 1 pint per 1,000 sq. ft. of soffits, mairing, window sills, outside of penned enclosures and on floor of food storage rooms.

ACCEPTED

12/16/71

U.S. FEDERAL INSECTICIDE AND RODENTICIDE ACT
REGISTERED UNDER NO. 912-31

COOP

WITH

DIRECTIONS (Cont)

Food Processing Milk Rooms and forms — For Use by Operators Only.

mosquitoes, wasps, silverfish and spider sprayer or fogger upward into the room. Apply 1 pint per 8,000 cu. ft. Do not apply space while food processing in operation. To control wasps, roaches, ants and silverfish, spray to cracks of and floors, behind and around door and window sills. Apply 1 pint per 1,000 sq. ft. Do not apply to food processing plants. For removed, and food has been covered with wrap application or will be effective. About 5 minutes after application, milking rooms will be

WITH 1% VAPONA[®]
INSECTICIDE

For Dairy Cattle, Beef Cattle, Food Processing Plants, Dairy Barns, Poultry Barns. Use in foggers or sprayers.


KILLS: Face flies, horn flies, stable flies, mosquitoes, gnats, roaches, ants, silverfish and sowbugs.

— FAST KNOCKDOWN OF INSECTS —

Active Ingredients — 100%


2,2-Dichlorovinyl Dimethyl Phosphate*	93%
Related Compounds*	.07%
Mineral Oil	30.00%
Petroleum Distillate	69.00%
(R) Trademark Shell Chemical Company	
* Equivalent to 1% Vapona Insecticide.	

WARNING: Keep Out of Reach of Children
See back panel for additional warning.



U.S.D.A. Registration 912-31

CO-OP



NET CONTENTS 1 GALLON

Manufactured by
FARMERS UNION



Central Exchange

INCORPORATED
ST. PAUL, MINN. 55101

Warning: May be inhaled or absorbed. Wash thoroughly with soap and water after handling and smoking. Do not eat, drink, or use tools or milking utensils. Do not induce vomiting. Get medical attention if inhaled. 2-PAM is a parasympatholytic. Do not administer with atropine. Do not use near open flame.

This product is toxic to fish. Do not use in or out of lakes, streams, ponds, or other water bodies. Do not contaminate water, soil, or equipment. Do not dispose of empty containers in water. Do not reuse containers.

OOP
VAPONA

COOP

CC

ACCEPTED
12/16/71
REGISTERED UNDER NO. 912-31

WITH 1% VAPONA[®]
INSECTICIDE

WITH 1% VAPONA[®]

DIRECTIONS (Continued)

Food Processing Plants, Dairies,
Milk Rooms and Outside Plat-
forms — For Use By Pest Control

Operators Only. To control flies, mosquitoes, wasps, roaches, ants, silverfish and spiders. Use hand sprayer or fogger and direct spray upward into the four corners of room. Apply 1 pint per 8,000 cu. ft. of space. Do not apply space spray treatments while food processing plant is in operation. To control wasps on nest, spray nest directly, at night. To control roaches, ants and silverfish, direct the spray to cracks of woodwork, walls and floors, behind shelving and under large equipment. Spray ant trails around door and window sills. Use 1 pint per 1,000 sq. ft. Respray as necessary. Do not apply this treatment in the edible products areas of food processing plants. Food should be removed, and food handling equipment covered with wrapping paper during application or washed afterward. About 5 minutes after spraying, ventilate rooms well before using.


For Dairy Cattle, Beef Cattle, Food Processing Plants, Dairy Barns, Poultry Barns. Use in foggers or sprayers.

KILLS: Face flies, horn flies, stable flies, mosquitoes, gnats, roaches, ants, silverfish and sowbugs.


— FAST KNOCKDOWN OF INSECTS —

Active Ingredients — 100%	
2,2-Dichlorovinyl Dimethyl Phosphate*	.93%
Related Compounds*	.07%
Mineral Oil	30.00%
Petroleum Distillate	69.00%
(R) Trademark Shell Chemical Company	
* Equivalent to 1% Vapona Insecticide.	

WARNING: Keep Out of Reach of Children
See back panel for additional warning.



U.S.D.A. Registration #12-31



NET CONTENTS 1 GALLON

Manufactured by
FARMERS UNION
Central Exchange
BY PAUL, MINN. 55101

Warning: May be fatal if swallowed, inhaled or absorbed through the skin. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate food-stuffs or milking utensils. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not spray on or near open flame.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not reuse container. Destroy when empty.

— To control
le flies and
fluid oz. per
daily. Spray
surfaces of
legs, but not
the animal's
id forehead)
tion against
er 2 fluid oz.



Beef Cattle
1 horn flies
arn fogger to
20. ft. Before
movement by
xs. Open and
minutes after
minute feed.
utensils. Do
ve received a
m within 8

y 1 pint per
manure, win-
ied enclosures
age rooms.

KILLS: Face flies, horn flies, house flies, mosquitoes, ticks, roaches, rats, silver fish and sowbugs.

FAST AND DOWN OF INSECTS

WARNING: Keep Out of Reach of Children
See back panel for additional warning.



NET CONTENTS 1 GALLON

U.S.D.A. Registration 912-31



Manufactured by
FARMERS UNION
Central Exchange
ST. PAUL, MINN. 55101

Warning: May be fatal if swallowed, inhaled or absorbed through the skin. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate food-stuffs or milking utensils. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not spray on or near open flame.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not reuse container. Destroy when empty.