

CO-OP[®] 20-20-12 SYSTEMIC - T
 Contains 0.5 THIMET[®] Soil Insecticide
 Fertilizer plus Insecticide

For use on Potatoes, Peas, Corn and Sugar Beets to control the insects listed below.

Active Ingredients:


Phosphate (Oxide) 15 - [(20) (10)]
 Phosphate (Sulfate) 6.8
 Percentally from fertilizer 100.0

Fertilizer Ingredients:

Total Nitrogen (N) 20
 Available Phosphorus (P₂O₅) 20
 Water Soluble Potash (K₂O) 12

DANGER!

KEEP AWAY FROM CHILDREN



WARNING:

Poisoning may occur as a result of exposure. May be fatal if swallowed, inhaled or absorbed through the skin. Do not breathe the dust. Avoid contact with skin, eyes or clothing. Wash all contaminated clothing with soap and hot water before reuse. Wash thoroughly after handling and before eating or smoking. Do not contaminate food or feed. Do not reuse empty container. Burn empty bags and stay away from smoke or fumes.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Apply this product only as specified on the label.

Directions

Thimet is an effective soil and systemic insecticide for early season insect control, which has been combined with fertilizer for efficient and uniform application. Application should be made at planting time in the row, to the side of the seed bed, not apply directly to the seed.

Recommended Application for Major Crops

<u>Crop</u>	<u>Insects</u>	<u>Rate - lbs per acre</u>	<u>Application</u>
Potatoes	Aphids, leafoppers, leafminer, psyllids, fireworms, flea beetle larvae, and reduction of flea beetle adults	250 to 375	Granules may be banded on each side of the row at planting time. Use 250 lbs on sandy or light soils, 375 lbs on heavy, firm, clay loam or truck soils. (30" row spacing)
Corn at planting time	Corn rootworms	63 to 125	At planting time, in the row - 1 1/2" or 1" above seed. Do not place in direct contact with seed. Use 125 lbs on fields having high rootworm populations.
Sugar Beets	Sugar Beet root maggot, leafhoppers, wire aphid, leafminer	125	Apply in 7" band over the row at planting time. Do not place in contact with seed. Make no application later than 30 days before harvest. Do not feed to livestock, poultry or other animals.
Beans	Leafhoppers, psyllids, fly, thrips, thrips, wireworm, rootworm	125 to 150	Distribute the granules in the row to the side of the seed bed. Apply 7-1 1/2" above the seed. Do not place in direct contact with seed. Do not feed to livestock, poultry or other animals.

©1 Trademark of Union Carbide Corporation
 ©2 Union Carbide Corporation, New York, N.Y.

Product of Union Carbide Corporation
 DURHAM, N.C. 27701
 Denver, Colorado

Distributed by
 Company
 State, City, and Zip

U.S. Dept. of Agriculture