

Net Weight 25 lbs.


Covers 5,000 sq ft.

GREENHOUSE

LAWN WEED & FEED contains 2,4-D and Silvex

DOES 3 BIG JOBS AT SAME TIME

- (1) KILLS MOST BROAD-LEAVED TYPE LAWN WEEDS--
DANDELION, BUCKHORN, PLANTAIN.
- (2) KILLS MOST VINING-TYPE LAWN WEEDS--
CHICKWEEDS, GROUND IVY, CLOVER.
- (3) FEEDS LAWN-CONTAINS SPECIAL REVITALIZER OF IRON AND
ADDITIONAL PLANT NUTRIENTS FOR EXTRA GREENING.



ACTIVE HERBICIDE INGREDIENTS

Dimethylamine Salts of 2,4-Dichlorophenoxyacetic Acid** 0.7%

Isooctyl Ester of Silvex***
((2-(2,4,5-Trichlorophenoxy) propionic acid)) 1.4%

INERT HERBICIDE INGREDIENTS 97.9%

TOTAL 100.0%

** Equivalent to 2,4-Dichlorophenoxyacetic Acid 0.00%

*** Equivalent to Silvex ((2-(2,4,5-Trichlorophenoxy)
propionic acid)) 0.00%

CAUTION: Keep out of reach of children.

See Additional Warnings and Directions on Back Panel.

12 point type

Manufactured by
CENEX, INC.
SOUTH ST. PAUL, MINNESOTA 55075

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 912-41

GREENHOUSE LAWN WEED AND FEED

GRASS FERTILIZER PLUS ADDITIONAL PLANT NUTRIENTS AND WEED CONTROL.

NET WEIGHT 25 LBS.

11-4-7

GUARANTEED ANALYSIS

Total nitrogen (N)	12%
3.5% Ammoniacal Nitrogen	
8.2% Water Soluble Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	8%
Additional Plant Nutrients	
Sulfur (S)	1.00%
Boron (B)	0.029%
Copper (Cu)	0.050%
Iron (Fe)	0.100%
Manganese (Mn)	0.050%
Molybdenum (Mo)	0.0005%
Zinc (Zn)	0.050%

Potential Acidity equivalent to 515 lbs. calcium carbonate per ton.

Primary plant foods derived from Urea, Phosphoric acid, and Muriate of Potash.

GREENHOUSE LAWN WEED AND FEED
 COVERS 5,000 SQUARE FEET

GREENHOUSE LAWN WEED AND FEED kills most broad-leaved and vining-type lawn weeds while it feeds your lawn. Kills Buckhorn, Dandelion, Curly Dock, Plantain, Thistles, Slickweed, Clover, Ground Ivy, and Parslane. 2,4-D and Silvex are combined with Lawn Food to provide feeding action and help grass thicken to fill in bare spots left by weeds.

DIRECTIONS FOR USE

WHEN TO APPLY

Use anytime weeds are actively growing and when temperature remains 50 degrees or above. For maximum effectiveness, apply on moist foliage, when lawn is damp from dew, rain or sprinkling. Control is slower if soil is dry or weather cool.

HOW TO APPLY

Apply over entire lawn area. Use a spreader set for an application rate of 5 lbs. per 1,000 Sq. Ft. Shut off spreader as you stop or turn around. Rain or sprinkling, mowing or heavy traffic within 24 hours reduces effectiveness, and a repeat treatment may be needed--wait at least one week. Delay planting new grass for two weeks. Do not apply to new plantings until after grass has been mowed at least twice. Grass may become discolored temporarily in periods of high heat or humidity, but with watering will make a fairly rapid recovery.

SPREADER SETTINGS

The settings for spreaders other than Co-op True Temper-Broadcase are only approximates. Age and condition of spreader will affect application. User should carefully observe rate of application at beginning of spread and make adjustment if necessary. The following settings are designed to apply 5 lbs. per 1,000 sq. ft.: Cyclone Broadcast Spreader, 6 ft. swath, setting of $4\frac{1}{4}$; Scotts 24" Model No. 75-3, setting of 9; Co-op True Temper Broadcast--10 ft. coverage, setting of $4\frac{1}{2}$.

CAUTION:

Avoid inhaling dust, avoid contact with skin, eyes or clothing. Avoid contamination of food or feedstuffs. Wash hands after using. **ANTIDOTE:** If taken internally, induce vomiting and call a physician immediately.

WARNING:

Use only when there is no hazard of drift. May injure bentgrass, St. Augustine, Dichondra, carpet grass, and grass seedlings in newly seeded ground. Do not apply directly to or otherwise permit the product to contact vegetables, flowers, grapes, fruit trees, ornamentals or other desirable plants which are sensitive to 2,4-D and similar materials. Excess amounts of 2,4-D acid in the soil may temporarily inhibit seed germination or plant growth. Do not store near fertilizers, seeds, insecticides or fungicides. Do not use on crops that will be utilized for food or forage purposes. Do not contaminate water used for domestic or irrigation purposes.

Thoroughly clean spreader equipment with a suitable cleaner before using for other purposes.

Vapors from this product may injure susceptible plants in the immediate vicinity.

Manufactured by
Cunox, Inc.
307th St. Paul, Minnesota 55105

