

Fertilizer plus insecticide

Contains 1.0% D-D-SYSTEM

Statement of Ingredients for Insect Control

Active Insecticide Ingredients:

O,O-Diethyl S-(2-(ethylthio) ethyl) phosphorodithioate 1.0%

Inert Insecticide Ingredients:

99.0%

100.000%

Statement of Fertilizer Ingredients

Total Nitrogen (N) 6%

Available Phosphoric Acid (P₂O₅) 26%

Water Soluble Potash (K₂O) 26%

WARNING: Keep Out of Reach of Children

Accidents may occur as a result of exposure. May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Avoid contact with skin, eyes, or clothing. Do not contaminate food or feed. **Wash thoroughly after handling.**

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Do not apply where run-off is likely to occur.

TO PHYSICIAN: Atropine Sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Directions for Use

D-D-SYSTEM is a systemic insecticide that enters the plants through the root system. Broadcast the granules near the roots, is picked up and moves in the xylem to the feeding parts of the plant where it kills insects as they feed on the plant.

Apply D-D-SYSTEM FERTILIZER plus INSECTICIDE as recommended herein, opposite the root system of the plant. A second treatment of granules may be necessary in some cases to insure adequate coverage and to control any insects which may be present at the time of application.

IMPORTANT: When used on plants which are susceptible to root rot, apply a fungicide to the soil around the plants to prevent root rot.

Do not use on plants which are susceptible to root rot. Do not use on plants which are susceptible to root rot.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Recommended Uses and Insects Controlled

Crop	Insect	Rate-lbs/acre	Remarks
ALFALFA	Aphids, mites	100	Apply at planting time. Allow a minimum of 28 days between applications.
BEANS (Snap-Lima-Dry)	Aphids, leafhopper, mites, thrips and Mexican bean beetle larvae	100 to 200	Apply at planting time in a broadcast or band application. Make only one application per season. At planting time, apply 100 lbs/acre. At second application, apply 200 lbs/acre. for feed, food or silage within 40 days of application.
CORN	Rootworms, aphids, leafhoppers, mites and thrips	100	Apply at planting time in a broadcast or band application. Do not apply within 75 days of harvest.
PEAS	Aphids, leafhoppers	200	Apply at planting time in a broadcast or band application. Do not apply within 75 days of harvest.
POPLARS	Aphids, flea beetle, leafhopper, psyllid, spruce sawfly, spruce budworm, spruce beetle, spruce sawfly, spruce budworm, spruce beetle	200 to 400	Apply at planting time in a broadcast or band application. Do not apply within 75 days of harvest.
SOYBEANS	Aphids, leafhoppers	100	Apply at planting time in a broadcast or band application. Do not apply within 75 days of harvest.

(North Dakota and Minnesota)

USDA Reg No 912-42

ACCEPTED
JAN 10 1969
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 912-42