

## DIRECTIONS

### ONE QUART MAKES 25 TO 240 GALLONS OF SPRAY OR DIP

**BEEF CATTLE AND HORSES:** For control of Lone Star Tick, Ear Tick, Cattle Fever Tick and Horn Flies, dilute one quart in 25 gallons of water. For Ear Ticks treat infested ears, neck and head. Dip or apply to animal thoroughly with enough spray force to part hair and wet the skin. Do not allow beef cattle access within 30 days of slaughter. Do not spray within 30 days of slaughter or dip within 60 days of slaughter. Do not apply to animals under 3 months of age.

**FOR LICE ON BEEF CATTLE:** Dilute 1 quart in 48 gallons of water. Thoroughly spray. If calves are infested and must be treated do not dip if they are under 6 months of age but apply spray sparingly or sponge on infested areas. Do not add infested animals to the treated herd. Do not spray within 30 days of slaughter or dip within 60 days of slaughter. Do not apply to animals under 3 months of age.

**SHEEP AND GOATS:** Fleece Worms - Sheep Scab - Lice and Sheep Ticks. For Fleece Worms mix 1 quart to 25 gallons of water. For Sheep Scab mix 1 quart with 48 gallons of water. For Lice and Sheep Ticks (Keds) mix 8 ounces with 60 gallons of water. Dip or spray thoroughly, preferably in Spring or Summer when wool is short to give insecticide spray penetration. Replacement Ewes or Rams should be treated before adding to the flock. For Sheep Scab, dip animals making sure that material penetrates wool, especially around head. If dipping vat is intermittently used, agitate and recharge to the desired strength before next dipping. Do not spray within 30 days nor dip within 60 days of slaughter. Do not use on Dairy Goats. Do not apply to animals under 3 months of age.

**HOGS:** For Lice mix 1 quart in 25 gallons of water. A single application will give control. Spray sows 30 to 45 days before farrowing. Do not treat pigs that are less than weaning age. Do not treat within 30 days of slaughter. Do not treat sows for 3 weeks after farrowing. Do not apply to animals under 3 months of age.

**SARCOPTIC MANGE MITE:** For use on Beef Cattle, Hogs, and Horses, mix 1 quart in 25 gallons of water. Apply as a dip or spray for thorough animal coverage. Repeat application in 15 days. Do not treat within 30 days of slaughter. Do not treat animals under 3 months of age. When dipping, intermittently agitate the mixture to maintain a milky color and obtain good consistent coverage.

CO-OP®

ACCEPTED  
9-9-71  
912-56  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR THE UNITED STATES OF AMERICA  
REGISTERED

# STOCK -TOX PLUS

LIVESTOCK SPRAY CONCENTRATE

for Beef Cattle, Hogs, Sheep and Horses

CONTROLS: LICE, TICKS, FLIES, MOSQUITOES,  
SARCOPTIC MANGE MITES, FLEAS

CO-OP

*Beef Cattle*

**BACKRUBBERS:** For control of Horn Flies and as an aid in reduction of Face Flies, dilute 1 quart with 11 quarts of fuel oil. Apply the diluted material to saturate the backrubber mechanism. Recharge the reservoir or treating surface as needed. For Horn Flies locate the backrubber to give the cattle free access to all parts of the body, particularly the back. For Face Flies one end of the backrubber must be low enough (not more than 15 to 20 inches above ground) to allow animals to treat their heads. Effective only for those cattle treating face and head. Specially designed canvas face treaters may be available in your area. Do not allow calves under 6 months of age to treat themselves.

Do not allow dairy cattle access to backrubbers. Do not allow beef cattle access within 30 days of slaughter. Do not allow animals under 3 months of age access to backrubbers. **UTILITY SPRAY for Mosquitoes, Flies, Fleas and Gnats—treat Out-Buildings, Loading Sheds, Barns, Fences and Waste Foliage,** dilute 1 quart to 25 gallons of water. Thoroughly spray to point of runoff. Make application to wall, doors, ground areas, ceilings and places where insects rest or breed. Do not use in Dairy Barns or Milk Houses. Do not spray foliage that may be eaten by domestic animals. Repeat applications as needed for control. For Fleas treat animal bedding and replace with new. Do not use in dwellings. Do not contaminate feed. Do not use in poultry houses. Do not use utility spray dilution on animals.

**WARNING:** May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or clothing. Wash thoroughly after using. Do not contaminate feed or food stuffs. In case of contact with eyes or skin wash immediately and get medical attention for eyes. May be absorbed through the skin.

Avoid treating overheated or sick animals. Avoid treating animals in cold stormy weather.

Do not apply this product to Dairy Animals. Do not apply in Dairy Barns, Milk Houses or Poultry Houses nor to pets or humans. This product cannot be used on food or forage to be fed to Dairy Animals or animals being finished for slaughter. Water animals before dipping.

Do not treat animals under 3 months of age. This product is toxic to fish and wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Keep out of any body of water. Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning (keep out of smoke).

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

