

**DIRECTIONS FOR USE**

**MIXING DIRECTIONS**—FOR HOSE SPRAYERS AND GARDEN PRESSURE SPRAYERS, MIX 2 TABLESPOONS PER GALLON OF WATER APPLIED. A SMALL AMOUNT OF LIQUID DETERGENT MAY BE MIXED WITH SPRAY TO HELP WET THE WEEDS.

**WEEDS KILLED**—Common and mouse-eared chickweed, chicory, clover, dandelion, dock, henbit, plantain, ragweed, thistle, and wild onion. Do not use on dichondra or lippia lawns. Make two applications for plants such as thistle, bindweed, wild onion, spotted spurge and knotweed or as needed for control.

**WHEN TO SPRAY**—For chickweed, treat in the fall before seeds are produced (October). Repeat if needed for complete kill. For other susceptible weeds listed, spray in late spring when weeds are producing new growth or as needed through the season. Reseeding can be done within 3 to 4 weeks. Do not spray new lawns until after second cutting.

**SPRAYING**—Apply 1 gallon of dilute spray to 180 sq. feet of turf. To get best results, area treatment spray preferred over spot spraying. Spray again if rain occurs before it has dried on weeds. Do not use on bent grass, buffalo, carpet grass or St. Augustine lawns since damage will result.

Packed for  
Farmers Union Central Exchange, Inc.  
St. Paul, Minnesota 55165



**LAWN WEED KILLER**  
WITH SILVEX and 2,4-D

KILLS CHICKWEED,  
DANDELION,  
PLANTAIN  
AND MANY  
OTHER  
WEEDS

8 OUNCES MAKES 8 GALLONS  
OF DILUTE SPRAY

**ACTIVE INGREDIENTS:**

\*Mixed octyl ester of 2,4-Di-chlorophenoxyacetic acid ..... 14.0%

\*\*Mixed octyl ester of Silvex 2-(2,4,5-Trichlorophenoxy propionic acid)..... 4.3%

**INERT INGREDIENTS** ..... 81.7%

Total ..... 100.0%

\*Equivalent to 9.2% 2,4-Dichlorophenoxy-acetic acid

\*\*Equivalent to 3.0% Silvex 2-(2,4,5-Trichlorophenoxy) propionic acid

**CAUTION: Keep Out of Reach of Children**  
See Side Panel for Additional Cautions

Contents: 1 Pint E.P.A. Reg. No. 912-57

ACCEPTED  
MAY 12 1971  
UNITED STATES DEPARTMENT OF AGRICULTURE  
FEDERAL AGENCY FOR REGULATORY AND COMPLIANCE

**DIRECTIONS FOR USE**

**WOODY PLANTS**—For poison ivy in fence row and vacant lots, mix 8 ounces in 2½ gallons of water. Use a garden pressure sprayer. Add 1 tablespoon of liquid detergent. Thoroughly wet ivy. Do not wet ivy climbing up tree trunks, spray sparingly and repeat as needed. Repeat 2 weeks later if needed. For stump resprouting, mix 8 ounces in 1 quart of kerosene. Soak cut surfaces, especially near bark and base of stump.

**CAUTION**—Avoid contact with skin, eyes and clothing. Harmful if swallowed. Do not use in greenhouses. Vapors from this product may injure susceptible plants in immediate vicinity. Do not store in confined spaces with living plants. Do not allow spray to drift to nearby susceptible plants such as grapes, garden vegetables, roses or other flowers. Coarse sprays are less likely to drift. Do not use hose applicators when near susceptible plant. Do not store with or near seeds, fertilizers, insecticides or fungicides. Do not contaminate water used for irrigation or domestic purposes.

This product will kill fish, keep out of any body of water.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Label Code No. 57-671

ACCEPTED WITH COMMENTS



**DIRECTIONS FOR USE**  
**MIXING DIRECTIONS—FOR HOSE SPRAYERS AND GARDEN PRESSURE SPRAYERS.** MIX 2 TABLESPOONS PER GALLON OF WATER APPLIED. A SMALL AMOUNT OF LIQUID DETERGENT MAY BE MIXED WITH SPRAY TO HELP WET THE WEEDS.

**WEEDS KILLED—**Common and spotted chickweed, chickory, dandelion, dock, henbit, plantain, ragwort, thistle, and wild onion. Do not use on dichondra or Hippia lawns. Make two applications for plants such as thistle, bindweed, wild onion, spotted spurge and knotweed or as needed for control.

**WHEN TO SPRAY—**For chickweed, treat in the fall before seeds are produced (October). Repeat if needed for complete kill. For other susceptible weeds listed, spray in late spring when weeds are producing new growth or as needed through the season. Reseeding can be done within 3 to 4 weeks. Do not spray new lawns until after second cutting.

**SPRAYING—**Apply 1 gallon of dilute spray to 100 sq. feet of turf. To get best results, area treatment spray preferred over spot spraying. Spray again if rain occurs before it has dried on weeds. Do not use on bent grass, buffalo, carpet grass or St. Augustine lawns since damage will result.

Packed for  
**Farmers Union Central Exchange, Inc.**  
 St. Paul, Minnesota 55165

**LAWN WEED KILLER**  
**WITH SILVEX and 2,4-D**

KILLS CHICKWEED,  
 DANDELION,  
 PLANTAIN  
 AND MANY  
 OTHER  
 WEEDS

8 OUNCES MAKES 8 GALLONS  
 OF DILUTE SPRAY

**ACTIVE INGREDIENTS:**

\*Mixed octyl ester of 2,4-D; chlorophenoxyacetic acid 14.0%

\*\*Mixed octyl ester of Silves 2-(2,4,5-trichlorophenoxy) propionic acid 4.3%

**INERT INGREDIENTS** 81.7%

Total 100.0%

\*Equivalent to 9.2% 2,4-Dichlorophenoxyacetic acid

\*\*Equivalent to 10% Silves 2-(2,4,5-trichlorophenoxy) propionic acid

**CAUTION: Keep Out of Reach of Children**

See Side Panel for Additional Cautions

Contents: 1 Pint E.P.A. Reg. No. 912-57

**ACCEPTED**  
 Jan 12, 1973  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC PEST CONTROL  
 UNDER NO. 912-57

**DIRECTIONS FOR USE**

**WOODY PLANTS—**For poison ivy, in fence row and vacant lots, mix 8 ounces in 1/2 gallons of water. Use a garden pressure sprayer. Add 1 tablespoon of liquid detergent. Thoroughly wet ivy. Do not wet ivy climbing up tree trunks, spray sparingly and repeat as needed. Repeat 2 weeks later if needed. For stump resprouting, mix 8 ounces in 1 quart of kerosene. Soak cut surfaces, especially near bark and base of stump.

**CAUTION—**Avoid contact with skin, eyes and clothing. Harmful if swallowed. Do not use in greenhouses. Vapors from this product may injure susceptible plants in immediate vicinity. Do not store in confined spaces with living plants. Do not allow spray to drift to nearby susceptible plants such as grapes, garden vegetables, roses or other flowers. Coarse sprays are less likely to drift. Do not use hose applicators when near susceptible plant. Do not store with or near seeds, fertilizers, insecticides or fungicides. Do not contaminate water used for irrigation or domestic purposes.

This product will kill fish, keep out of any body of water.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Label Code No. 57-671