

# 10% <sup>BRAND</sup> Chlordane WETTABLE POWDER

## INSECTICIDE

**Controls: Ants, Chiggers, White Grubs, Wire Worms, Flies**

|                       |               |
|-----------------------|---------------|
| ACTIVE INGREDIENTS    |               |
| Technical Chlordane*  | 10.00%        |
| Petroleum Hydrocarbon | 2.38%         |
| Inert Ingredients     | <u>87.62%</u> |
|                       | 100.00%       |


\*Contains 6% Octachloro 4, 7-dimethanotetrahydroindane and 4% related compounds.)

**CAUTION: Keep Out of Reach of Children**

See Side Panel for Additional Cautions

E.P.A. Reg. No. 912-59

NET WEIGHT ONE POUND



Packed for  
**Farmers Union Central Exchange, Inc.**  
So. St. Paul, Minnesota 55165

## DIRECTIONS FOR USE

**TO USE AS A DUST:** Dust from sifter top lid or garden duster. **IN LAWNS** to control Ants, Chiggers and Chinch Bugs use 1/2 lb. per 1,000 square feet. Repeat as needed for control. For White Grubs and Sod Webworms apply 9 oz. per 500 square feet. For Ants, Grubs and Sod Webworms water dust in. For Chiggers and Chinch Bugs leave dust on foliage. **IN CABBAGE** for Cut Worms and Mole Crickets on Lettuce, Pepper and Beets dust on soil and plants with 1/3 lb. per 1,000 square feet. Repeat within 10 days or as needed. Soil treatments should be limited to preplanting, planting, or transplanting time. Do not apply after edible part start to form on peppers and lettuce.

**AS A SPRAY IN LAWNS:** Use 15 to 22 tablespoons in 1 gallon water to cover 300 square feet. Apply in the spring to control White Grubs, White Fringed Beetle larvae, Ants, Chiggers, Sod Webworms and Chinch Bugs. For Ants treat hills and trails. Repeat application as needed for control. Agitate spray solution frequently. Birds feeding on treated areas may be killed. Keep children and pets off treated areas until this product is washed into the soil and the grass is dry.

**CRABGRASS:** Apply in the early spring before crabgrass germinates. Use 15 lbs. to cover 1,000 square feet. Use fertilizer spreader and water dust down to the soil level. This application will also kill Lawn Grubs, Webworms and Ants. Keep children and pets off treated areas until this material has been thoroughly washed into the soil and the grass has completely dried. For use on lawns. Do not use on newly seeded lawns.

**FRUITS:** When spraying make sure that solution is agitated often. To control Plum Curculio on apples and peaches mix 4 tablespoons in 1 gallon water. On apples make first application at petal fall and repeat at 7 to 10 day intervals. On peaches apply at shuck stage or first cover. Repeat if necessary. Do not apply to apples or peaches within 30 days of harvest. To control Strawberry Crown Borer apply 1/3 lb. in 3 gallons water to 1,000 square feet. Spray plants just before the first blossoms appear. Do not apply after fruit begins to form.

**BEANS:** To control Cabbage Worms, Flea Beetles, Green Stink Bugs, Harlequin Bug and Mole Crickets apply 3/4 lb. per 1,000 square feet. Use as preplanting, planting, and transplanting soil treatment. For these insects on Cauliflower apply in the same manner but not after the curd begins to grow.

**CUCUMBER AND SQUASH:** To control Cucumber Beetles and Squash Bugs apply 5 1/2 oz. dust per 1,000 square feet or spray 11 tablespoons in 1 gallon to 500 square feet. Do not apply after edible portions begin to form. To prevent plant injury, do not apply while plants are wet with dew.

**LETTUCE:** For Cutworms, Lygus Bugs, and Serpentine Leafminer apply as for Bean insects. Do not apply after heads begin to form or to leafy lettuce after the seedling stage.

**ONIONS:** For Onion Maggot use at same rate as for Bean insects. Apply as a preplanting, planting or transplanting soil treatment.

**POTATOES:** For Blister Beetle control apply to foliage at the rate of 9 tablespoons in 1 gallon water for 500 square feet. For White Grubs in the soil treat the same as for Grubs in the Lawn.

**RESIDUAL PAINT OR SPRAY FOR FLIES, MOSQUITOES, GNATS, ROACHES, AND WATERBUGS:** Mix 4 oz. (16 tbsps.) in 1 1/4 pint water. Apply by painting or spraying to walls and areas where insects rest or run in farm buildings, utility animal sheds, porches and basements. Do not apply in dairy barns or milk houses. One gallon should cover about 1,000 sq. ft. Spray mixture should be frequently agitated to keep particles in suspension. Spray directly as many insects as possible directing spray into cracks and crevices especially for Roaches and Waterbugs. Use only as a spot spray in homes. Keep out of reach of children. Do not use in poultry houses. Do not use in homes for fly control.

**CAUTION:** Do not breathe dust or spray mist. Avoid contamination of feed and foodstuffs. Wash thoroughly after use. Keep out of reach of children.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label. Apply only on calm days to avoid dust or spray drift.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.