



Tomato and Garden Dust

DIRECTIONS FOR USE

Start application as soon as plants are set or immediately after true leaves form. Repeat every 7 to 10 days and after every rain. Follow the caution statements as regards latest treatment possible to avoid residues at harvest. Apply dust to cover both sides of leaves and the stems as well as possible. Dust on wet or dry foliage but preferably when air is still. **TOMATO and GARDEN DUST** will control insects and plant diseases listed below:

BEANS: Aphids, Leafhoppers, Mexican Bean Beetle, Bean Leaf Beetle, Red Spider, Thrips, Japanese Beetle, Cucumber Beetle, and Downy Mildew.

BEETS and CARROTS: Leafhoppers, Blister Beetles, Aphids, Early and Late Blight and Leaf Spot. (If tops are to be used for food and feed, do not apply to Beets and Carrots within 14 days of harvest.)

CANTALOUPEs, WATERMELONS and CUCUMBERS: Melon Worm, Striped Cucumber Beetle, Spotted Cucumber Beetle, Flea Beetle, Melon Aphid, Spider Mite, Leaf Spot and Downy Mildew or Blight.

PEPPERS: Spinach Aphids, Anthracnose and Cercospora.

POTATOES: Early Blight, Colorado Potato Beetle, Potato Aphid, Leafhopper, Potato Thrips, Potato Flea Beetle, Potato Psyllid, False Chinch Bug and Mealybugs.

CABBAGE: Aphids, Imported Cabbage Worm, Cabbage Looper and Diamondback Moth.

LETTUCE: Leafhoppers, Aphids, Mites and Cabbage Lopper. (Do not apply to Leaf Lettuce within 14 days of harvest.)

SWEET CORN: Corn Earworm, Aphids, Sap Beetles, Thrips (Injury may occur if applied in the whorl stage.) Do not apply within 14 days of harvest.

PUMPKINS: Aphids, Mites, Leafhoppers and Squash Vine Borers.

SQUASH: Alfalfa Loopers, Aphids, Spider Mites, Pickleworm, Squash Vine Borer, Cucumber Beetles (Do not apply unless plants are dry.)

PEAS: Pea Aphid, Pea Weevil, Alfalfa Looper and Celery Looper.

RADISHES: Aphids

TOMATOES: Spider Mites, Aphids, Tomato Russet Mites, Flea Beetles, Leafhoppers, Psyllid, Drosophila, Leaf Spot, Anthracnose, Early and Late Blight.

ROSES: Leaf Spot, Aphids and Rose Chafer (Surface of Bed should also be dusted).

FLOWER INSECTS: Chrysanthemum Thrips, Tussock Moth, Japanese Beetle, Thrips, Thrips on Snapdragons, White Fly and Sowbugs.

CAUTION: Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Remove residues remaining at harvest by washing, wiping, or similar effective means. Do not use on ferns or crassula. Do not apply to any of the listed crops within 7 days of harvest unless otherwise specified.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.



This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

Packed for:

Farmers Union Central Exchange, Inc.
So. St. Paul, Minnesota 55165

Label Code No. 60-



Tomato and Garden Dust



CONTROLS INSECTS and DISEASES

ACTIVE INGREDIENTS:

*Methoxychlor Technical	5.0%
**Captan	5.0%
***Malathion	4.0%
INERT INGREDIENTS	86.0%
TOTAL	100.0%

*Equivalent to 4.4% 2, 2-bis (p-methoxyphenyl) - 1, 1, 1-trichloroethane and 0.6% other isomers and reaction products.

**N ((trichloromethyl)thio) - 4-cyclohexene - 1, 2 - dicarboximide.

***O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

NET CONTENTS

EPA Reg. No. 912-60

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See other cautions on back panel