

CAPITOL

... CONCENTRATE

ACTIVE INGREDIENTS

PYRETHRINS-----.75%
*PIPERONYL BUTOXIDE
technical-----3.75%
DEODORIZED PETROLEUM
HYDROCARBONS---95.50%
100.00%

*Equivalent to 3.00% of (Butyl Carbityl)
(5-tropyk Piperonyl) Ether and 0.75% of
related compounds.

908-26


DIRECTIONS: Use in accordance with recommendations
of Marketing Research Report #334, by Joseph N.
Tenhet, Stored-Tobacco Insects Laboratory, Richmond,
Virginia.

CIGARETTE BEETLE: Applied at the rate of 3 fluid
ounces per 1,000 cubic feet of air space, (in a fine
mist). Repeat as necessary.

HAZIFUL IF SWALLOWED. DO NOT SPRAY, POUR OR SPILL NEAR
OPEN FLAME. AVOID INHALATION OR SPRAY OR MIST.

NET CONTENTS 55 GAL DRUM

USDA Reg. # 107-



CAPITOL CHEMICAL COMPANY
Sanitation Consultants

450 BUTLER ROAD
WASHINGTON, D.C. 20006

[212-61]

CO-OP[®]
LIVESTOCK
Spray

ACCEPTED

MAY-19-1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
REGISTERED PERSON REGISTERED

912-61

INSECTICIDE

Approved for use on dairy animals

CO-OP
LIVESTOCK
spray

DIRECTIONS

Apply as a mist over dairy cows, cattle, horses, and hogs. Do not wet animal's hide, only a mist is required. Do not apply spray in poultry houses.

To develop a residual repellency to stable flies, horn flies, mosquitoes and gnats on dairy cows, apply 1 to 2 ounces of LIVESTOCK SPRAY to each animal for a three day period. The spray should be applied with a good sprayer so that the entire animal's body has been uniformly covered. Special care should be taken in spraying the legs, belly and shoulders. Avoid areas around the nose, eyes and mouth. Spray in the same manner again after two days have elapsed, and every second day thereafter or when insects again disturb the animal. Be sure the teats and udder are washed with warm water prior to milking. Calves should be sprayed very lightly once a day instead of attempting to build up a residual repellency by spraying more heavily.

For control of lice on cattle, horses, and hogs, spray the animals with 2 ounces of LIVESTOCK SPRAY. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks. To kill flies, mosquitoes and gnats in the barn prior to milking, close all doors and windows.