

# CAUTION


Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing spray mist or vapor. In case of contact with skin wash with soap and water. Avoid contamination of feed and foodstuffs. Do not use or store near heat or open flame.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain.

Clean application equipment and dispose of containers, rinse solutions, wastes, and unused chemical by burying in non-cropland away from water supplies.

Use only in compliance with the limitations, cautions and warnings stated on this label.

Keep children and pets off treated grass and turf until the insecticide has been washed off the grass and into the soil, either by adequate rainfall or thorough sprinkling, and the grass has completely dried. Do not feed or graze grass from treated areas to livestock or poultry. Do not use grass from treated areas for hay. Birds feeding over treated lawn or ornamental turf may be harmed.



BRAND

# CHLORDANE 4EC

CONTROLS MAJOR INSECTS IN LAWNS AND GARDENS  
INSECTICIDE

**ACTIVE INGREDIENTS:**

\*Technical Chlordane.....42.37%

Aromatic Petroleum Hydrocarbons .....53.63%

INERT INGREDIENTS .....4.00%

TOTAL 100.00%


Equivalent to 25.42% Octachloro-4, 7-methanotetrahydroindane and 16.95% related compounds (Contains 4 lbs. Chlordane per Gallon)

**CAUTION: Keep Out of Reach of Children**  
See Side Panel for Additional Cautions

EPA REG NO 912-66

NET CONTENTS 1 QUART

Packed for: Farmers Union Central Exchange  
So. St. Paul, Minnesota 55165




**DIRECTIONS FOR USE Dilute in sufficient water**

CROP	INSECTS	RATE/1,000
SWEET CORN	Corn rootworm(Northern and Southern) Cornfield ants, Corn root aphids	2 4 - 5 1/4 tbsp.
	Seed corn beetle larvae, Seed corn maggots, Cutworms, White grubs, Wireworms, White-fringed beetle larvae, Asiatic garden beetle larvae	3 1/2 - 5 3/4 tbsp.
VEGETABLES: Beans, Beets, Broccoli, Brussel sprouts, Cabbage, Cantaloupe, Cauliflower, Celery, Collards, Cucumbers, Eggplant, Kale, Kohlrabi, Lettuce, Melons, Okra, Onions, Peas, Peppers, Pimentos, Potatoes, Radishes, Rutabagas, Squash, Tomatoes, Turnips, Watermelon	Cutworms, Wireworms, White grubs	3 2 - 5 3/4 tbsp.
	Asiatic garden beetle larvae, Whitegringed beetle larvae, Flea beetle larvae(Potatoes)	3 2 - 7 1/3 tb
	Wireworms (potatoes-Pacific Northwest)	1/4 - 1/2 pt.
FRUITS AND BERRIES: Apples, Cherries, Nectarines, Pears, Plums, Prunes, Quinces, Blackberries, Grapes, Huckleberries, Loganberries, Raspberries, Strawberries, Kumquats,	Ants	2 4 - 2 3/4 tbsp.
	White grubs, Cutworms, Wireworms	3 2 - 5 3/4 tbsp.
	Asiatic garden beetle larvae, Whitefringed beetle larvae	3 2 - 7 1/3 tb
	Blackberries, Blueberries, Raspberries, Strawberries	1/2 pt.
LAWNS and ORNAMENTAL TURF	Rootworms	1/4 - 1/2 pt.
	Chiggers, Ants, Ticks	2 4 tbsp.
	Earthworms	1/2 - 2 1/3 pts.
AROUND BUILDINGS and HOUSES	White grubs, Japanese beetle grubs, Crickets, Sod Webworms	3 2 - 7 1/3 tb
	Ants, Earwigs, Roaches, Sowbugs, Spiders, Centipedes, Crickets, Silverfish	5 1/2 tbsp. in 1 water

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT**

Accepted With Comments



BRAND

# CHLORDANE 4EC

CONTROLS MAJOR INSECTS IN LAWNS AND GARDENS  
INSECTICIDE

**ACTIVE INGREDIENTS:**

\*Technical Chlordane.....42.37%  
Aromatic Petroleum Hydrocarbons .....53.63%

**INERT INGREDIENTS** .....4.00%

**TOTAL 100.00%**

Equivalent to 25.42% Octachloro-4, 7-methanotetrahydroindane and 16.95% related compounds (Contains 4 lbs. Chlordane per Gallon)

**CAUTION:** Keep Out of Reach of Children  
See Side Panel for Additional Cautions

EPA REG NO 912-66

NET CONTENTS 1 QUART

Packed for: Farmers Union Central Exchange  
So. St. Paul, Minnesota 55165

**REGISTERED**  
FEDERAL INSECTICIDE AND RODENTICIDE ACT  
MICROBIC REGISTERED

**DIRECTIONS FOR USE** Dilute in sufficient water to cover the area indicated

CROP	INSECTS	RATE/1,000 sq. ft.	DIRECTIONS
SWEET CORN	Corn rootworm(Northern and Southern) Cornfield ants, Corn root aphids	2 4 - 5 3/4 tbsp.	On Sweet Corn and Vegetables, make preplant application to soil after plowing and work into soil immediately.
	Seed corn beetle larvae, Seed corn maggots, Cutworms, White grubs, Wireworms, White-fringed beetle larvae, Asiatic garden beetle larvae	3 1/2 - 5 3/4 tbsp.	
VEGETABLES: Beans, Beets, Broccoli, Brussel sprouts, Cabbage, Cantaloupe, Cauliflower, Celery, Collards, Cucumbers, Eggplant, Kale, Kohlrabi, Lettuce, Melons, Okra, Onions, Peas, Peppers, Pimentos, Potatoes, Radishes, Rutabagas, Squash, Tomatoes, Turnips, Watermelon	Cutworms, Wireworms, White grubs	3 2 - 5 3/4 tbsp.	
	Asiatic garden beetle larvae, Whitefringed beetle larvae, Flea beetle larvae(Potatoes)	3 1/2 - 7 1/3 tbsp.	
	Wireworms (potatoes-Pacific Northwest)	1/4 - 1/2 pt.	
FRUITS AND BERRIES: Apples, Cherries, Nectarines, Pears, Plums, Prunes, Quinces, Blackberries, Grapes, Huckleberries, Loganberries, Raspberries, Strawberries, Kumquats, Blackberries, Blueberries, Raspberries, Strawberries	Ants	2 1/4 - 2 3/4 tbsp.	Make application and work into soil. On fruit and berries, do not apply when fruit is present.
	White grubs, Cutworms, Wireworms	3 1/2 - 5 3/4 tbsp.	
	Asiatic garden beetle larvae, Whitefringed beetle larvae	3 1/2 - 7 1/3 tbsp.	On berries, make application at base of plants and work into soil.
	Root weevil larvae	1/4 pt.	
	Rootworms	1/4 - 1/2 pt.	
LAWNS and ORNAMENTAL TURF	Chiggers, Ants, Ticks	2 4 tbsp.	Apply spray thoroughly to grass and repeat as necessary.
	Earthworms	1/2 - 2 1/3 pts.	
	White grubs, Japanese beetle grubs, Crickets, Sod Webworms	3 1/2 - 7 1/3 tbsp.	Apply to grass in early spring or late fall.
AROUND BUILDINGS and HOUSES	Ants, Earwigs, Roaches, Sowbugs, Spiders, Centipedes, Crickets, Silverfish	5 2 tbsp. in 1 gal. water	Apply to ground and foundations around buildings.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME**

Label Code No. 66-1072

**5% GRANULAR**

# **5% GRANULAR KILLS**

**ANTS, CUTWORMS, CHIGGERS, TICKS, LAWN  
MOTHS (SOD WEBWORMS), JAPANESE BEETLE  
LARVAE, EUROPEAN CHAFER LARVAE.**

**ACTIVE INGREDIENTS**

Technical Chlordane*	5.0%
INERT INGREDIENTS	95.0%
<b>TOTAL</b>	<b>100.0%</b>

\*Equivalent to 3% of Octachloro-4,7-methanotetrahydroindane and 2% of related compounds.

**CAUTION:** KEEP OUT OF REACH OF CHILDREN. For additional cautions see back panel.

E.P.A. Reg. No. 961-153  
E.P.A. Est. 961-PA-1

**MANUFACTURED BY  
LEBANON CHEMICAL CORPORATION**

LEBANON, PA. 17040

# — DIRECTIONS —

Granular 5% Chlordane broadcasts easily in a fertilizer Spreader settles quickly to the soil and will not dust. The following settings may be used as a guide to achieve the application rates presented below. All settings are approximate and every spreader should be carefully calibrated on a measured plot. Factors such as variations in the walking rate, age of spreader, etc.

Spreader	1 lb 1,000 sq. ft.	2 lb 1,000 sq. ft.	4 lb 1,000 sq. ft.
Wagon	3 1/2	4 1/2	5
MTD	4	4 3/4	5 1/2
Queen Turt	4	4 3/4	5 1/2
Central	4	5	5 1/2
Seas	4	5	5 1/2
Jackson	4	4 3/4	5 1/2
Glanor	4	5	5 1/2
Cyclone	2 3/4	3	3 1/2

## FOR SURFACE INSECTS

**ANTS** Apply Granular 5% Chlordane to soil surface and individual mounds at the rate of 1 lb. per 1,000 sq. ft.

**CUT WORMS, CHIGGERS, TICKS** Treat lawn surface with 1 lb. Granular 5% Chlordane per 1,000 sq. ft.

**CATERPILLARS, MOTHS, AND WEB WORMS** Apply 2 lbs. Granular 5% Chlordane per 1,000 sq. ft. in the late evening. Water with 1 gallon of water per 1,000 sq. ft.

**NOTE:** Keep children and animals off treated areas until the material has been thoroughly washed into the soil and the treated area is completely dry. Do not graze treated area.

## FOR SOIL INSECTS

**ANTS** Apply 1 lb. Granular 5% Chlordane per 1,000 sq. ft.

**SOIL WEB WORMS, LAWN MOTH (ANNA)** Water lawn thoroughly after application. Apply 2 lbs. Granular 5% Chlordane per 1,000 sq. ft. in the late evening.