

Directions

To control certain plant diseases use this product at the rate of 2½ tablespoons per gallon of water or 2 lbs. per 100 gallons, unless otherwise indicated.

Apples and Pears — Apple scab, fruit rots, and blotch — make applications at 7 to 10-day intervals as needed to maintain cover in pre-bloom, bloom, petal fall and successive cover sprays. Continue applications to within 30 days of harvest to control blotch, up to start of harvest to control fruit rots. Do not use on D'Anjou pears. Russetting may be produced on Bosc pears.

Cherries, Peaches, Plums and Prunes — Brown rot, cherry leaf spot, blossom blight, peach scab — apply at pink bud, full bloom and petal fall stages. Repeat at 10 to 14-day intervals until harvest.

Raspberries, Blackberries, Dewberries — Anthracnose — begin application at first bloom. Make two repeat applications at 10 to 15-day intervals.
Botrytis fruit rot — Apply 3 to 5 days before harvest and repeat in mid-harvest and again 8 to 16 days later.

Strawberries — Botrytis fruit rot and leaf spot — make application when new growth starts and repeat at 7-day intervals through harvest.

Cantaloupes, Cucumbers, Pumpkins, Summer & Winter Squash, Watermelons — Angular leaf spot, Anthracnose, downy mildew. Apply at first sign of disease and repeat at 5 to 7-day intervals or as needed to maintain disease control.

Asparagus Preplanting Root Dip — (For Damping-off Control) — Use 3 lbs. per 100 gals. of water (4 tablespoonfuls per gal.) Dip root sections for 1 minute, drain and plant.


Tomatoes — Anthracnose, early and late blight, gray leaf spot, septoria leaf spot — Apply at first sign of disease. Repeat at 5 to 7-day intervals as needed to maintain control. Use 3 tablespoons per gallon of water except where blight conditions are severe. Under these conditions increase dosage to 5 tablespoons per gallon of water.

Greenhouse Captan 50W BRAND

A fungicide for plant disease control

Active Ingredients:

Captan (N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	50%
Inert Ingredients:	50%
Total	100%



CAUTION: Keep out of reach of children
Avoid inhalation of dust or spray mist.
Avoid contact with skin. Avoid contamination of feed and foodstuffs.


This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Containers should be disposed of by burying with wastes, or by burning (keep out of smoke). Do not contaminate water by cleaning of equipment, or disposal of wastes.

Net Contents

Packed for
FARMERS UNION CENTRAL EXCHANGE, INC.,
P.O. Box G, St. Paul, Minnesota 55165

912-73



Carrots — Septoria leaf spot, Cercospora blight — Use 5 tablespoons per 2 gallons water. Apply at first sign of disease and repeat at 6 to 10-day intervals

Potatoes — Early and late blight — Timing and rates of application same as tomatoes above.

Soil Treatment — Preplanting for damping-off and root rot diseases on seedlings or transplants of Roses (and other shrubs, trees, flowers), Lawn Seedbeds, Vegetables (such as beans, peas, celery, crucifers, eggplant, pepper, tomato) — Use 4 to 5 ozs. per 1,000 sq. ft. as a dust, or 2 lbs. per 100 gals. water at a rate of 15 gals. per 1,000 sq. ft. Cultivate into upper 3 to 4 inches of soil before planting.

Lawns & Turf — Brown patch, damping-off, leaf spot, root rot, melting out — Use 2 lbs. per 100 gallons or 2½ tablespoons per gallon and apply to surface at the rate of 10 gallons per 1,000 sq. ft. of turf. Repeat at 7 to 10-day intervals as needed. Apply at first signs of disease. For melting out, apply when grass resumes growth in the spring. Repeat at 7 to 10-day intervals as long as cool, wet weather persists. Do not graze treated areas. Do not feed clippings to livestock.

Roses — Black Spot — Apply at first signs of disease. Repeat at 7 to 10-day intervals, more frequently during frequent rains and heavy dews.

Preplant Dip for Gladiolus, Tuberous Begonias — Seed rot, damping off — Mix 6 to 8 tablespoons per gallon of water. Immerse bulbs in mixture for 20 to 30 minutes. Drain and plant.

Carnations, Chrysanthemums — Rust, leaf spot, flower blight — Apply at first signs of disease. Repeat applications as needed at 7 to 10-day intervals.

Note: Always follow good sanitation practices. Destroy or burn all diseased vegetation.

E.P.A. Reg. No. 912-73

LCN: 73-473