

Front Panel

Hooker Agricultural Chemicals

MBC[®]

Granular, Nonselective Grass and Weedkiller

Apply Dry Or As A Spray For

General Vegetation, Perennial Noxious Weeds, Preparing Treatment

USDA Registration No. 935-15

Active Ingredients

Sodium Chlorate (NaClO₃) 30%

Sodium Metaborate Tetrahydrate (Na₂B₂O₄·4H₂O) 68%

(Boron Trioxide (B₂O₃) Equivalent 23.3%)

Inert Ingredients 2%

CAUTION

Keep out of reach of children.

Harmful if swallowed.

Do not contaminate feed and food stuffs.


Do not contaminate water used for domestic purposes or irrigation ditches.

Avoid contact with skin and clothing.

Wash clothing and equipment after application.

Do not apply to desirable plants or areas underlaid by their roots.

ACCEPTED
APR 11 1967
UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT AS FUNGICIDE AND ROENTICIDE FOR ECONOMIC POISON REGISTERED UNDER NO. 935-15 SUBJECT TO ATTACHED COMMENTS



Keep animals off treated areas until rainfall has leached

the herbicide into the soil.

Mechanical equipment should be thoroughly cleaned after use.

Do not store in direct contact with ground or concrete floors

IMPORTANT

See Reverse Side for Directions for Use and Other Precautions

LB NET


HOKER CHEMICAL CORPORATION

Niagara Falls, New York Columbus, Mississippi

Manufactured under U.S. Patent No. 2,800,500

Back Panel

MBC[®] GRANULAR



APPLICATION For all uses, uniform application is essential. MBC Granular may be applied either as a dry granular material or as a water spray. Dissolve in water at rates up to 3 pounds per gallon. Both spray and dry applications become effective as soil treatments following rainfall or irrigation.

For DRY application, conventional herbicide spreaders may be used.

For Spray application, equipment with mechanical agitation is recommended.

Add up to 3 pounds MBC Granular per gallon of water, depending on job

requirements, to spray tank. Continue agitation until the herbicide

is completely dissolved and, preferably, until spray tank is empty.

To minimize fire hazard due to dry vegetation, treatment while weeds

are green is suggested.

GENERAL VEGETATION CONTROL - MBC Granular will give quick knock down

of vegetation and residual control of broad leaved weeds and grasses.

ACCEPTED
 Apr 11 1967

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR REGISTERED USE ONLY
 TO ATTENTION

935-15

on parking lots, fence lines, ^{sites} highway and railroad sites, drainage
 ditches and other similar areas. For SPRAY applications, add 1/2
 to 2 pounds MBC Granular to sufficient water to cover 100 square feet
 and apply as a contact spray, thoroughly wetting all parts of the
 plants, particularly the growing tips -- 1/2 to 1 gallon per 100 square
 feet is usually adequate for young growth up to 6 inches high. The
 higher rates may be necessary to insure thorough coverage of heavy
 vegetation or to provide extended residual control. For DRY granular
 treatment, apply 1/2 to 2 pounds per 100 square feet. The lighter
 rates should be used only at time of weed emergence on moist soil or
 when precipitation is anticipated while weeds are still young.

Persistent perennial weeds are most effectively treated when nearing
 maturity or during dormant stages in the fall. Spring applications may
 also be made under moderate rainfall conditions. Summer applications
 are recommended only when rainfall is dependable for leaching the
 herbicide into the soil. Apply 2 to 4 pounds per 100 square feet in

ACCEPTED
apr 11 1967
UNDER THE FEDERAL INSECTICIDE
REGULATIONS AND PESTICIDE ACT
FOR ECONOMIC POISON REGISTERED
EPA REG. NO. 935-15 SUBJECT
T-SUB-48 COMMENTS

the dry granular form or as a spray. On deep-rooted perennials

bindweed, Canada thistle, Bermuda grass and Quack grass, the heavier

rates may be required.

JOHNSON GRASS CONTROL - MBC Granular is particularly effective on

deep-rooted perennial grasses such as Johnson grass and Dallis grass

on industrial sites, road sides, fence lines and drainage ditches.

Apply dry at 1 to 3 pounds per 100 square feet or add 1 to 3 pounds to

1 to 2 gallons of water to adequately cover 100 square feet. Maximum

efficiency on Johnson grass is obtained by treating after heading to

late fall.

FOR USE UNDER ASPHALT PAVING - For use as a "prepaving" residual

herbicide under asphalt paving to prevent premature breakdown of the

paving due to weeds, an application of 2 to 4 pounds per 100 square feet

to the finished subgrade is recommended.

Apr 11 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTS REGISTERED
ED UNDER N. 935-15 FIRST
TO ATTACHED COMMENTS

For SPRAY treatment, dissolve 1 to 2 pounds per gallon of water and
apply 2 gallons per 100 square feet.

For DRY treatment, apply in granular form. After treatment, apply at
least 2 gallons of water per 100 square feet to dissolve and incorporate
the herbicide in the soil.

NOTICE

This material is not selective in action and may be toxic to all
vegetation and may render the entire treated area totally or partially
unproductive for one or more years. Care should be taken to confine
the use or application to the particular area intended to be treated
and to avoid its contact with lawns, trees, shrubs, crops and other
desirable plants which are not intended to be destroyed or injured.
This includes precautions in treating areas which may be overlaid by
roots of adjacent valuable growths. Careless application of this
material or washing by water run-off to areas where desirable plants
are growing or which will be used for later plantings may result in
injury to desirable plants. No responsibility is assumed by manufacturer

ACCEPTED
APR 11 1967
UNDER THE FEDERAL INSECTICIDE
HERBICIDE AND MOLLUSCIDANT
ACT FOR ECONOMIC PESTS CONTROL
REGISTRATION NO. 135-15 SUBJECT
TO ATTACHED COMMENTS.

or seller for unproductivity of the soil, injury to plant life or other results of this material when the above instructions are not followed. Because of the many varying conditions affecting use and application such as location, seasonal weather, soil composition and type of vegetation, manufacturer makes no representation or warranty with respect to results or effects of such use whether or not used or applied in accordance with directions herein contained. Neither manufacturer nor seller makes, and their employees and agents are not permitted to make any representations or warranties, express or implied by law, not specifically set forth on the label. The buyer accepts this material subject to these terms and assumes the risk of usage and handling. It is advisable to consult state agricultural experiment station weed specialists or the extension service on specific problems.

ACCEPTED
Apr 11 1967
UNDER THE
FUNDS
935-10

HOOVER CHEMICAL CORPORATION

Manufactured under U.S. Patent No. 2,700,604

Niagara Falls, New York

-

Columbus, Mississippi