

ACCEPTED  
6/11/69  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 954-4



# POISON ANTIDOTE



If swallowed, give a tablespoonful of table salt in a glass of warm water and repeat until vomit fluid is clear. Give milk or white of egg beaten with water. **CALL A PHYSICIAN IMMEDIATELY.**

## WARNING!

**CAUSES IRRITATION OF SKIN, EYES, NOSE AND THROAT.** Do not breathe vapor. Use with adequate ventilation. Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush skin or eyes with plenty of water; for eyes, get medical attention.

Printed in U.S.A. Lot 667 373

# FORMALTABS®

RELEASES  
**FORMALDEHYDE GAS**  
FOR DRY CABINETS

Active Ingredient:  
Formaldehyde (polymeric) 93%  
Inert Ingredients 7%

KEEP BOTTLE COVERED



# POISON



**DANGER! KEEP OUT OF REACH OF CHILDREN. SEE LEFT PANEL FOR ANTIDOTE STATEMENT AND OTHER WARNING STATEMENTS.**

KING RESEARCH, INC., Brooklyn, N.Y. 11215, Distr.

## SANITARY STORAGE CABINETS IN BEAUTY AND BARBER SHOPS Directions for Use

FORMALTABS release formaldehyde gas upon exposure to air in cabinet. Use 2 FORMALTABS for each 1/2 cubic foot of enclosed air space. (Use 3 tablets for the standard shop cabinet.)

Place FORMALTABS in small open dish and set on shelf of cabinet. Keep door closed at all times. **Do not mix the FORMALTABS with water.**

FORMALTABS slowly disappear as they release formaldehyde gas. Add fresh FORMALTABS periodically to replace material thus converted into gas.

**Net Weight: 3.52 Oz. (100 grams)**  
(50 2 gram tablets)  
**U.S.D.A. Reg. No. 954-4**