

UNDER THE FEDERAL
FUNGICIDE AND RODENT ACT
FOR ECONOMIC INSECTICIDES
REGISTERED UNDER NO. 160-123 SUBJECT
TO ATTACHED COMMENTS.

SMALL FRUIT VEGETABLES (FRUITING PLANTS)

SMALL FRUIT VEGETABLES (FRUITING PLANTS)
FIELD CROPS - AMOUNTS RECOMMENDED ARE
AMOUNTS TO USE PER ACRE IN SUFFICIENT
WATER TO COVER

~~BLACKBERRY~~ ~~BOYSBERRY~~ ~~DEARBERRY~~
~~LOGANBERRY~~ ~~RASPBERRY~~
~~CRANBERRY~~

~~GRAPES (3) Slender Mince/Mediterranean~~
~~STRAWBERRY (3)~~

~~BEANS (1)~~
~~BEETS (7)~~ ~~BRUSSELS SPROUTS (7)~~

~~CABBAGE (7)~~ ~~KALE (7)~~ ~~MUSTARD (7)~~ ~~CAULIFLOWER~~
~~(7)~~ ~~BROCCOLI (3)~~ ~~TURNIP (3)~~

~~CELERY (7)~~
~~CUCUMBER (7)~~ ~~MELON (7)~~ ~~SQUASH (7)~~

~~EGGPLANT (3)~~

~~LETTUCE (1)~~

~~ONIONS~~

~~Peanut~~

~~PEPPERS (3)~~ ~~POTATOES~~

~~TOMATOES~~

~~WATERMELON~~

~~ZUCCHINI~~

Apply Malathion Premium Grade to the surface of plants
by directed spray at the rate of 1.0 to 1.5 gallons of water
per acre.

DIRECTIONS FOR CROP USE

Malathion 55 Insecticide Premium Grade

CONTAINS
CYTHION

THE PREMIUM GRADE MALATHION

CAUTION

KEEP OUT OF REACH OF CHILDREN

CATTLE, BEEF AND NON MILKING
Apply Malathion Premium Grade to the surface of plants by directed spray at the rate of 1.0 to 1.5 gallons of water per acre.

DOGS
Apply Malathion Premium Grade to the surface of plants by directed spray at the rate of 1.0 to 1.5 gallons of water per acre.

SHEEP, GOATS
Apply Malathion Premium Grade to the surface of plants by directed spray at the rate of 1.0 to 1.5 gallons of water per acre.

POULTRY
Apply Malathion Premium Grade to the surface of plants by directed spray at the rate of 1.0 to 1.5 gallons of water per acre.

PILGRIM TIT
Apply Malathion Premium Grade to the surface of plants by directed spray at the rate of 1.0 to 1.5 gallons of water per acre.

