


Best Available Copy



DIRECTIONS FOR USE

IMPORTANT: Read these Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using ZIPCIDE Cattle Dust Bag.

CONDITIONS OF SALE: These Directions for Use reflect the opinion based on tests of effectiveness, of toxicity to plants and laboratory animals, and of residue on the food and feed normal conditions of use. However, because of the wide range of conditions under which the product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions. The control of horn flies, BALCOM CATTLE DUST BAG, and the safety of the animal are the subject of this statement. The user assumes all

HORN FLY

Suspend ZIPCIDE CATTLE DUST BAGS in areas frequented by cattle or in places through which the animals pass daily. ZIPCIDE Cattle Dust Bags may also be suspended in the shade or in front of protected areas for protecting dairy cows. ZIPCIDE Cattle Dust Bags may be suspended in the exit through which the milking herd leaves the milking parlor. In all cases, the dust bags should be adjusted so that the bottom of the bag is about six inches below the top of the animal's head.

CATTLE DUST BAG

* Contains * CO-RAL (or coumaphos)

FOR CONTROL OF HORN FLIES ON BEEF AND DAIRY CATTLE

GUARANTEE

ACTIVE INGREDIENT:

O, O-Diethyl O-(3-chloro-4-methyl-2-oxo-2H-1,4-benzopyran-7-yl) phosphorothioate 1%

INERT INGREDIENTS: 99%

U.S. Patent No. 2,748,146;
Canadian Patent No. 523,206
U.S.D.A. Registration No. 960-169

CAUTION

KEEP OUT OF REACH OF CHILDREN
See right panel for Caution statements

NET WEIGHT 10 POUNDS

*CO-RAL is Reg. TM in U.S. and Canadian Pat. Offs. by Faberfabrik Bayer and Chemagro Corp. Licensee.

INSTRUCTIONS FOR USING CATTLE DUST BAG

Suspend the dust bag by hanging rope, chain, or wire through the grommet at the top of the bag and attaching to an overhead structure or by hooking the grommet into hooks fixed to a stationary overhead and at the bottom of the bag, attach the two pins and hang the bag in the center.

Install as many 10 pound ZIPCIDE Cattle Dust Bags as necessary in lanes or other openings to assure per cent of the animals. When dust bags are used on a large dairy herd, allow one 10 ZIPCIDE Cattle Dust Bag for 10 to 20 animals. The bags should be placed in all lanes and insecticide should be applied from the bottom.

RESTRICTIONS: Do not hang cattle dust bags near feed, drinking water troughs.

CAUTION: Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and water after handling.

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, consult prompt medical aid. Prolonged exposure will result in dermatitis or depression.

To Physician—Atropine sulfate is antidote. Atropine is also antidotal and may be administered in conjunction with atropine.

BALCOM CHEMICALS, INC.

240 22 ND. STREET P.O. BOX 637 GREELEY, COLORADO 80631 303 - 352-4750

THREAD AND PULL HARD TO ZIP OPEN BOTTOM

TYPE OF CROWNEE
EVENLY GUIDING THE KNOT IN ROPE

BE

960-169 8-11-1969