





# CATTLE DUST BAG

## DAIRY

### DIRECTIONS FOR USE

**IMPORTANT:** Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using ZIPCIDE CATTLE DUST BAGS.

**CONDITIONS OF SALE:** These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. BALCOM CHEMICALS, INC. and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

### LOUSE AND HORN FLY CONTROL:

Suspend ZIPCIDE CATTLE DUST BAG in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or mineral. Dust bags may also be suspended in loafing sheds or in front of protected mineral feeders. For Lactating dairy cows, ZIPCIDE CATTLE DUST BAGS may be suspended in the exit through which cows leave the milking barn. In all cases, the dust bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle.

### LOUSE AND HORN FLY CONTROL AND AS AN AID IN REDUCING FACE FLY POPULATIONS:

Suspend ZIPCIDE CATTLE DUST BAG in the same manner as described under "Louse and Horn Fly Control" except as follows: (1) The bags should be hung low enough to insure that the faces of the cattle will be dusted. (2) The bags must be located so the animals will be forced to use them daily. As face flies are able to travel over an area of several miles, best

CATTLE D  
CONTAINS  
(or cou

FOR CONTROL OF  
FLIES AND AS AN  
FACE FLY POPUL  
AND DAIR

GUAR  
ACTIVE INGREDIE  
0,0-Diethyl 0-(3-chloro  
-1-benzopyran-7-yl)  
INERT INGREDIEN

TO

U.S. Patent N  
Canadian Paten

EPA REG. N

CAUTION

KEEP OUT OF REA  
See right panel for

NET WEIGHT

\*CO-RAL is a Reg. T  
Bayer



# BALCOM CHEMICALS

240 22 ND. STREET P. O. BOX 667 GREENE



GRAB THREAD AND PULL HARD TO  
ZIP OPEN BOTTOM.

IT SLIDING TIE KNOT IN ROPE  
OF GROMMET

TO PREVENT SLIDING TIE KNOT IN ROPE  
THIS SIDE OF GROMMET




# BALCOM

# ZIPCIDE

®

DISPENSING BAG U.S. PATENT NO. 3,494,333

## CATTLE DUST BAG



# DAIRY

# BEEF

### DIRECTIONS FOR USE

**IMPORTANT:** Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using ZIPCIDE CATTLE DUST BAGS.

**CONDITIONS OF SALE:** These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. BALCOM CHEMICALS, INC. and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

**LOUSE AND HORN FLY CONTROL:**

### CATTLE DUST BAG CONTAINS CO-RAL\* (or coumaphos)

FOR CONTROL OF LICE AND HORN  
FLIES AND AS AN AID IN REDUCING  
FACE FLY POPULATIONS ON BEEF  
AND DAIRY CATTLE.

### GUARANTEE

**ACTIVE INGREDIENT:**  
0,0-Diethyl 0-(3-chloro-4-methyl-2-oxo-(2H)-  
1-benzopyran-7-yl) phosphorothioate... 1%


results in reducing face fly populations are achieved when ZIPCIDE CATTLE DUST BAGS are used on an area-wide basis.

### INSTALLATION OF ZIPCIDE CATTLE DUST BAG

Suspend the dust bags by inserting rope, chain, or wire through the grommets at the top of the bag and attaching to an overhead structure, or by hooking the grommets into hooks fixed to a stationary overhead 2x4 at the heights recommended for louse and horn fly or face fly control. Rope, chain or wire can also be tied or attached to two posts or other fixed points with the dust bag hanging in the center. When dust bags are provided on a free-choice basis, allow one 10 pound ZIPCIDE CATTLE DUST BAG per 10 to 20 animals. After the bags are in place, pull the rip cord to allow the insecticide to release from the sifter bottom.

**RESTRICTION:** Do not hang Cattle Dust Bag over feed, mineral, or water troughs.

This product is toxic to fish, birds, and other wildlife.



**DAIRY**



**BEEF**

**DIRECTIONS FOR USE**

**IMPORTANT:** Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using ZIPCID CATTLE DUST BAGS.

**CONDITIONS OF SALE:** These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. BALCOM CHEMICALS, INC. and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

**LOUSE AND HORN FLY CONTROL:**

Suspend ZIPCID CATTLE DUST BAG in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or mineral. Dust bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, ZIPCID CATTLE DUST BAGS may be suspended in the exit through which cows leave the milking barn. In all cases, the dust bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle.

**LOUSE AND HORN FLY CONTROL AND AS AN AID IN REDUCING FACE FLY POPULATIONS:**

Suspend ZIPCID CATTLE DUST BAG in the same manner as described under "Louse and Horn Fly Control" except as follows: (1) The bags should be hung low enough to insure that the faces of the cattle will be dusted. (2) The bags must be located so the animals will be forced to use them daily. As face flies are able to travel over an area of several miles, best

**CATTLE DUST BAG  
CONTAINS CO-RAL\*  
(or coumaphos)**

**FOR CONTROL OF LICE AND HORN  
FLIES AND AS AN AID IN REDUCING  
FACE FLY POPULATIONS ON BEEF  
AND DAIRY CATTLE.**

**GUARANTEE**

**ACTIVE INGREDIENT:**  
0,0-Diethyl 0-(3-chloro-4-methyl-2-oxo-(2H)-  
-1-benzopyran-7-yl) phosphorothioate ... 1%  
**INERT INGREDIENTS** ..... 99%  
**TOTAL** ..... 100%

U.S. Patent No. 2,748,145;  
Canadian Patent No. 523,206

EPA REG. NO. - 960-100

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**  
See right panel for caution statements.

**NET WEIGHT 10 POUNDS**

\*CO-RAL is a Reg. TM of Farbenfabriken  
Bayer Ag.

results in reducing face fly populations are achieved when ZIPCID CATTLE DUST BAGS are used on an area-wide basis.

**INSTALLATION OF ZIPCID CATTLE DUST BAG**

Suspend the dust bags by inserting rope, chain, or wire through the grommets at the top of the bag and attaching to an overhead structure, or by hooking the grommets into hooks fixed to a stationary overhead 2x4 at the heights recommended for louse and horn fly or face fly control. Rope, chain or wire can also be tied or attached to two posts or other fixed points with the dust bag hanging in the center. When dust bags are provided on a free-choice basis, allow one 10 pound ZIPCID CATTLE DUST BAG per 10 to 20 animals. After the bags are in place, pull the rip cord to allow the insecticide to release from the sifter bottom.

**RESTRICTION:** Do not hang Cattle Dust Bag over feed, mineral, or water troughs.

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not contaminate food or feed. Destroy damaged or worn out dust bag by burying or burning. Stay away from smoke or fumes. Do not breathe dust. Wash after handling. If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

**TO PHYSICIAN:**

Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish, birds, and other wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**WARRANTY AND LIMITATION OF DAMAGES**

BALCOM CHEMICALS, INC. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the risks referred to therein. BALCOM CHEMICALS, INC. makes no other expressed or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of BALCOM CHEMICALS, INC. is authorized to do so except in writing with specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.

**BALCOM CHEMICALS, INC.**

240 22 ND. STREET P. O. BOX 667 GREELEY, COLORADO 80631 303 - 352-4750



**B THREAD AND PULL HARD TO  
OPEN BOTTOM.**

**GRAB THREAD AND PULL HARD TO  
ZIP OPEN BOTTOM.**

