

CAPTAN 7.5 DUST

FUNGICIDE

FOR ECONOMIC POISONING
 ED UNDER NO. 960-189
NET WEIGHT
50 LBS.

DIRECTIONS FOR USE

- ① **APPLES:** Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Pox, Botryosphaeria — Use 25 to 40 lbs. per acre. Apply at delayed dormant period. Repeat at weekly intervals as required to maintain control.
- ② **APRICOTS:** Brown Rot, Jacket Rot — Use 50 to 60 lbs. per acre. Apply at red bud stage and repeat at 75% petal-fall.
- ③ **CHERRIES:** Cherry Leaf Spot, Brown Rot — Use 50 to 60 lbs. per acre. Apply at popcorn, blossom and petal fall stages. Continue at 10 to 14 day intervals as required.
- ④ **PEACHES:** Brown Rot, Peach Scab — Use 50 to 60 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal-fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary.
- ⑤ **PEARS:** Pear Scab (except Pacific Northwest) — Use 25 to 40 lbs. per acre. Apply during early finger and petal-fall stages. Repeat at weekly intervals as necessary to maintain control. Russeting may be reduced on Bosc pears. Do not use on D'Anjou pears.
- ⑥ **RASPBERRIES:** Fruit Rot — Use 30 lbs. per acre making application 3 to 5 days before harvest starts, at mid-harvest and 8 to 10 days later.
- ⑦ **STRAWBERRIES:** Botrytis Rot — Use 40 lbs. per acre. Make the first application when new growth starts in the spring and before the fruit starts to form. Repeat at weekly intervals. During severe infection periods continue through harvest, treating immediately after each picking.

ACTIVE INGREDIENT:

*Captan 7.5%

INERT INGREDIENTS: 92.5%

100.0%

*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide

U. S. Patents No. 2,553,770; 2,553,771; 2,553,776

CAUTION KEEP OUT OF REACH OF CHILDREN.

Avoid inhalation of dust. Avoid Contact with skin.
 Avoid Contamination of feed and foodstuffs.

USAGE CAUTION:

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke

EPA Reg. No. 960-189

⑧ **TOMATOES:** Anthracnose, Early Blight, Late Blight, Septoria Leaf Spot — On Young plants use 25 to 30 lbs. per acre. On mature plants use 40 to 50 lbs. per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

⑨ **POTATOES (SEED PIECES):** Seed Rot, Damping Off, Verticillium Wilt, Brown Eye Disease — Dust at the rate of 10 ounces per bushel of cut seed pieces (1 lb. per 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 to 3 days in open crates before bagging. Do not use treated seed for food or feed purposes.

Young Plants: Early Blight, Late Blight — Use 30 to 35 lbs. per acre. Begin applications when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens.

Mature Plants: Early Blight, Late Blight — Use 40 lbs. per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

⑩ **ORNAMENTALS, (ROSES, CARNATIONS, CHRYSANTHEMUMS):** Black Spot, Leaf Spot, Rust Flower Blight — Apply at first sign of disease. Repeat at 7 to 10 day intervals as necessary. Dust plants thoroughly.

NOTICE: Buyer assumes all risk of use, storage, or handling of this product when not in strict accordance with directions given herewith.



BALCOM CHEMICALS, INC.
 240 22ND STREET P O BOX 667 GREELEY COLORADO 80631 PHONE 303-352-4750

