MALATHION BIG-6 DUST

Und r the Federal Insecticie in Fungithie, and Rodenticide Act, as amenued, for the posticide registered under 2000.

CAUTION

May be harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after handling. Avoid contamination of feed and food stuffs.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

IMPORTANT

Use of this product, as directed will be within the tolerance of 8 parts per million for Malathion. TREATED GRAIN IS NOT SUBJECT TO DOWN-GRADING and can be milled into flour or feed using normal production methods.

NOTICE: Buyer assumes all risk of use, storage, or handling of this product when not in strict accordance with directions given herewith.

FOR USE ON STORED GRAINS: Wheat, Corn, Oats, Rice, Rye, Barley, Grain Sorghum, Field and Garden Seeds.

PROTECTS AGAINST DAMAGE FROM: Confused Flour Beetle, Rice Weevil, Granary Weevil, Saw-toothed Grain Beetle, Flat Grain Beetle, Red Flour Beetle, Rusty Grain Beetle, Lesser Grain Borer, Indian Meal Moth.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Side Panels for Additional Precautions

ACTIVE INGREDIENTS: BY WEIGHT

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate)

INERT INGREDIENTS:

100.0%

94.0%

6.0%

EPA Reg. No. 960-202

TOTAL

NET WEIGHT

LBS.

(PCLEAN CROP is Reg. T.M. of Balcom Chemical, Inc.

DIRECTIONS FOR USE

STORED GRAIN AND SEED: Apply MALATHION BIG 6 DUST to the grain or seed at the rate of 10 lbs. per 1000 bushels as the grain is being loaded or turned into final storage. To protect stored grain or seed from attack by Indian Meal Moth, apply this product to the surface of grain or seed at the rate of 5 lbs. per 1000 sq. feet of surface area. Apply the dust evenly over the surface immediately after the grain or seed is loaded into storage. Repeat if necessary.

HOW TO APPLY

One convenient way to apply MALATHION BIG-6 DUST to shelled corn and small grains is to treat it when it is in the field wagon or truck prior to binning. Determine the amount in the wagon or truck by multiplying the length by width by depth (all in feet) and multiply this tot al by 0.8 to get the number of bushels. Measure out the correct amount of MALATHION BIG-6 DUST and spread evenly over the surface of the grain in the wagon or truck bed. Cut the dust into the grain with a shovel. The dust will become mixed with the grain as it is hauled from the field to the storage bin, and as it is dumped from the bed into the auger hopper. Further mixing will occur when the grain is augered or blown into the storage bin. Mixing is completed as the treated grain rolls down the coneshaped pile in the bin. Another method of applying MALATHION BIG-6 DUST to grain or seed is to evenly meter the correct amount of dust onto the grain or seed as it is being transferred into storage, using a suitable metering device. Mixing is completed as the grain or seed is transferred into storage and rolls down the cone-shaped pile in the bin.

