


NET WT. 20 LBS.

COVERS 5,000 SQ. FT.

\$595



3/22/71

961-212

Comments

LEBANON

WEED AND FEED

10-6-4

WITH

SILVEX AND 2,4-D

CAUTION: KEEP OUT OF REACH OF CHILDREN. See back panel for additional cautions.

ACTIVE INGREDIENTS:	
Dimethylamine 2,4-dichlorophenoxyacetate*	.64 %
Isooctyl ester of Silvex 2-(2,4,5-trichlorophenoxy) propionic acid**	.75 %
INERT INGREDIENTS	98.61 %
TOTAL	100.00 %

*Equivalent to .53% 2,4-dichlorophenoxyacetic acid

**Equivalent to 53% Silvex 2-(2,4,5-trichlorophenoxy) propionic acid

U S D A Reg. No. 961-212

LEBANON CHEMICAL CORPORATION
LEBANON, PA. 17042

COVERS 5,000 SQ. FT.



3/22/71

961-212

WEED AND FEED

10-6-4 WITH SILVEX AND 2,4-D

MINIMUM GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N) -----	10.0%
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅) -----	6.0%
SOLUBLE POTASH (K ₂ O) -----	4.0%

Potential acidity 700 lbs. Calcium Carbonate per ton,

LEBANON CHEMICAL CORPORATION

LEBANON, PA. 17042

FOR A LUXURIOUS, WEED FREE LAWN

The 10-6-4 fertilizer stimulates grass growth to quickly conceal spots where weeds have died. Grows rich green grass and kills weeds at the same time.

EFFECTIVE BROADLEAF WEED CONTROL

Kills broadleaf weeds such as Dandelion, Buckhorn, Plantain, Chickweed, Clover, Ground Ivy, Henbit, Black Medic, Cinquefoil, Knotweed, Poison Ivy and Thistle.

EFFECTIVE BROADLEAF WEED CONTROL

Kills broadleaf weeds such as Dandelion, Buckhorn, Plantain, Chickweed, Clover, Ground Ivy, Henbit, Black Medic, Cinqufoil, Knotweed, Poison Ivy and Thistle.

SAFE FOR ESTABLISHED LAWNS

Will not harm most lawn grasses. Some grasses may brown slightly, but will recover quickly, especially if watered well when browning is noticed.

SPREADER SETTINGS

Lawn Beauty _____	5	Central _____	6	Sears _____	6
MTD _____	5	Queen Turf _____	5	Jackson _____	6
Cyclone _____	4	Scott _____	4½	Glorion _____	6

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 6 oz. per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the spreader settings necessary to apply the correct amount.

When applying with a spinner type spreader, use extreme caution not to apply too close to desirable plants.

DIRECTIONS

The best time to destroy weeds is when they are young before they crowd out good grasses. Apply when weeds are in active growth, at rate of 4 lbs. per 1,000 sq. ft. on established lawn.

Apply in Spring and early Fall when ground moisture is adequate for better weed control. During dry weather, water lawn thoroughly 24 hours prior to application.

For persistent weeds, when weeds reappear or if a heavy rain within 2 days after application, a second application may be necessary.

Treat new lawns only after 2nd or 3rd mowing.

Do not apply to creeping grass such as Bent Grass.

Do not mow, water or make heavy use of lawn for at least 24 hours after application.

CAUTION

Avoid inhaling dust. Avoid contact with skin, eyes or clothing.

Vapors from this product may injure susceptible plants in the immediate vicinity. Bentgrass, St. Augustine grass, Dichondra, Carpetgrass, or clover may be injured. Avoid drift to susceptible plants as this product may injure flowers, ornamentals and vegetables.

Do not store near fertilizer, seeds, insecticides or fungicides. Clean spreader thoroughly after use.

Do not reuse container. Destroy when empty.

WARRANTY

Lebanon Chemical Corporation warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ACCEPTED WITH COMMENTS