


**WITH  
2,4-D AND  
SILVEX**



NET WT. 8 LBS.

**\$5.45**

**COVERS  
5,000 SQ. FT.  
100' x 50'**

TO OPEN -  
FILTER TAPE  
THREAD.



**TURF**

**WEEPER MULCH**

# WEED KILLER



DANDELION



CHICKWEED



CLOVER



BUCKHORN

## KILLS



PLANTAIN



HENBIT



GROUND IVY



KNOTWEED

**CAUTION: KEEP OUT OF REACH OF CHILDREN (SEE BACK PANEL FOR ADDITIONAL CAUTIONS)**

**ACTIVE INGREDIENTS:**

Dimethylamine 2,4-dichlorophenoxyacetate*	1.32%
Isooctyl ester of Silvex (2-(2,4,5-trichlorophenoxy) propionic acid)**	1.05%
<b>INERT INGREDIENTS:</b>	97.63%
<b>TOTAL</b>	<b>100.00%</b>

\*Equivalent to 1.10% 2,4-dichlorophenoxyacetic acid

\*\*Equivalent to .75% Silvex (2-(2,4,5-trichlorophenoxy) propionic acid)


U.S.D.A. Reg. No. 961-213

## LEBANON CHEMICAL CORPORATION

LEBANON, PA. 17042

**8 LBS. COVERS 5000 SQ. FT.**

# **LEBANON TURF WEED KILLER SILVEX and 2,4-D**



## **FOR A WEED-FREE LAWN**

Kills broadleaf weeds such as Chickweed, Knotweed, Clover, Dandelion, Buckhorn, Plantain, Cinquefoil, Ground-Ivy, Henbit, Black Medic, Poison Ivy and Thistle.

## **APPLY WHEN WEEDS ARE IN ACTIVE GROWTH**

Set your spreader, fill and apply. No mixing. Cover entire lawn or spot-treat where weeds exist.

## **SPREADER SETTINGS**

LAWN BEAUTY .....	4½	JACKSON .....	6	GLORION .....	4
SCOTT .....	6	SEARS .....	5	QUEEN TURF .....	5
MTD .....	5	CENTRAL .....	4		

**NOTE:** Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply only 2½ ounces per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

## **DIRECTIONS**

LEBANON TURF WEED KILLER is a mixture of 2,4-D and Silvex for application to active growing broadleaf weeds. Better results are obtained if applied when ground moisture is adequate. During dry weather water lawn thoroughly 24 hours prior to application. For persistent weeds or if heavy rain within 2 days after application, a second application may be necessary.

## **FOR ESTABLISHED LAWNS**

With a calibrated spreader, apply evenly at rate of 1½ lbs. per 1000 sq. ft. Do not apply to creeping grass such as Bent grass. Do not mow, water or make heavy use of lawn for at least 24 hours after spreading.

## **FOR NEW LAWNS**

Do not apply to newly seeded lawn until after 2nd or 3rd mowing.

## **CAUTION**

Avoid inhaling dust. Avoid contact with skin, eyes or clothing. Vapors from this product may injure susceptible plants in the immediate vicinity. Bentgrass, St Augustine grass, Dichondra, Carpetgrass, or Clover may be injured. Avoid drift to susceptible plants as this product may injure flowers, ornamentals, etc.

