

NET WEIGHT 8 POUNDS
TREATS 2500 SQ. FT.

\$450



Glorion®

Glorion®

**LAWN WEED
AND
CRABGRASS
KILLER**

LAWN WEED AND CRABGRASS KILLER

ACCEPTED
3/1/71
INSECTICIDE
ANTICIDE AND
FUNGICIDE REGISTER
961-231



KILLS CRABGRASS PLANTS, DANDELION, BUCKHORN, PLANTAIN, CLOVER, HENBIT, GROUND IVY, CHICKWEED AND MANY OTHER BROADLEAF WEEDS.

CAUTION: KEEP OUT OF REACH OF CHILDREN. For additional cautions see back panel.

ACTIVE INGREDIENTS:	
Dicofol (Methoxyfenozide)*	54%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid**	4%
Isopropyl ester of Silver (2,4,5-trichlorophenoxy)acetic acid***	4%
INERT INGREDIENTS	38%
TOTAL	100%

*Total Aromatic, expressed as elemental arsenic, all in water soluble form 0.625%
**Equivalent to 0.667% 2,4-Dichlorophenoxyacetic acid
***Equivalent to 0.667% Silver (2,4,5-trichlorophenoxy)acetic acid
U.S.D.A. Reg. No. 241,231

MANUFACTURED FOR
NASSAU GARDEN, INC.
Lawn & Garden Division, Jericho, L. I., N. Y. 11753

WEED AND CRABGRASS KILLER

KILLER

6/10/71
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND WEEDICIDE ACT
 FOR THE ECONOMIC PEACON REGISTERED
 UNDER NO. 961-231

Comments



CAUTION: KEEP OUT OF REACH OF CHILDREN. For additional cautions see back panel.

ACTIVE INGREDIENTS:	
Dicofol Methenarsonate*	1.540%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid**	0.805%
Isocetyl ester of Silvex (2-(2,4,5-trichlorophenoxy) propionic acid)***	0.945%
INERT INGREDIENTS:	96.710%
TOTAL	100.000%
*Total Arsenic, expressed as elemental arsenic, all in water soluble form 0.625%	
**Equivalent to 0.667% 2,4-Dichlorophenoxyacetic acid	
***Equivalent to 0.667% Silvex (2-(2,4,5-trichlorophenoxy) propionic acid).	
U. S. D. A. Reg. No. 961-231	


MANUFACTURED FOR
NASSAU GARDEN, INC.
 Lawn & Garden Division, Jericho, L. I., N. Y. 11753

LAWN WEED AND CRABGRASS KILLER

**NET WEIGHT 8 POUNDS
TREATS 2500 SQ. FT.**

Glorion®

**LAWN WEED
AND
CRABGRASS
KILLER**



Comments

SPREADER SETTINGS

Lawn Beauty	6	Jackson	7
Central	6	Sears	7
Modern	7	Scott	6

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to 5 ounces per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

DIRECTIONS

Apply at first sign of crabgrass or weeds and anytime through the growing period when crabgrass or weeds are visible. Also controls immature goose grass, dallis grass, and yellow foxtail if applied when in seedling stage.

Apply at the rate of 3 lb. 2 oz. per 1,000 sq. ft. when weeds are in active growth. On new seeding delay application until after third mowing. Desirable grasses may discolor slightly but watering will make fairly rapid recovery. Clover, Bent grass and some other creeping grasses may be injured by this product.

GLORION®

LAWN WEED AND CRABGRASS KILLER

FROM THE FERTON INSECTICIDE
DIVISION AND NUTRITION
DEPARTMENT
ED. NUMBER MS 9127-231

Comments

KILLER

SPREADER SETTINGS

Lawn Beauty	6	Jackson	7
Central	6	Sears	7
Modern	7	Scott	6

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to 5 ounces per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

DIRECTIONS

Apply at first sign of crabgrass or weeds and anytime through the growing period when crabgrass or weeds are visible. Also controls immature goose grass, dallis grass, and yellow foxtail if applied when in seedling stage.

Apply at the rate of 3 lb. 2 oz. per 1,000 sq. ft. when weeds are in active growth. On new seeding delay application until after third mowing. Desirable grasses may discolor slightly but watering will make fairly rapid recovery. Clover, Bent grass and some other creeping grasses may be injured by this product.

For crabgrass apply when seedlings are visible at rate of 2 pounds per 1,000 sq. ft. on lawn which is wet from sprinkling or rain. Two applications at weekly intervals is recommended. Repeat schedule of application will be necessary when new crabgrass appears later in season.

Rain, mowing, watering or heavy traffic within 24 hours after application will reduce the effectiveness.

CAUTION

Harmful if swallowed. Avoid drift to susceptible plants as this product may injure flowers, ornamentals and vegetables. Avoid inhaling dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid contamination of feed or foodstuffs. Keep children, pets and livestock off treated areas. Do not store near fertilizer, seeds, insecticides or fungicides. Vapors from this product may injure susceptible plants in the immediate vicinity.

WARRANTY

Nassau Garden, Inc., warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

FNZS527-13B

14474SR5

FREIGHT SHIPPING BAG
Meeting requirements of
APPLICABLE FREIGHT CLASSIFICATION
Guaranteed by ST. REGIS PAPER CO.

13x2½x31, 4-130-FK

ACCEPTED WITH COMMENTS