



LEBANON GREEN GOLD 10-6-4

WITH CRABGRASS CONTROL

A
761-248
761-248

GUARANTEED MINIMUM FERTILIZER ANALYSIS

TOTAL NITROGEN N 10.0%
AVAILABLE PHOSPHORUS ACID P₂O₅ 6.0%
SOLUBLE POTASH K₂O 4.0%
Potential acidity 600 lb. calcium carbonate per ton.

LEBANON CHEMICAL CORPORATION

LEBANON, PA. 17042

Feeds your lawn kills crabgrass upon germination. Grass seed can be applied 60 days after applying GREEN GOLD WITH CRABGRASS CONTROL.

ESTABLISHES A WEED CONTROL ZONE

GREEN GOLD WITH CRABGRASS CONTROL establishes a weed control zone by killing certain annual weed seeds as they germinate. Application prior to weed germination will prevent poa annua, crabgrass, goosegrass, watergrass, yellow and green foxtail.

DIRECTIONS FOR USE

GREEN GOLD WITH CRABGRASS CONTROL can be used on a variety of turf or mixture of turf grasses.

Bluegrasses, ryegrass, fescue, bermudagrass, zoysiagrass, St. Augustinegrass, centipede-grass and bahiagrass are tolerant. Do not apply to newly sprigged areas of St. Augustine, Bermuda, or Centipede grasses.

SPREADER SETTINGS

Lawn Beauty.....	7	Central.....	5½	Modern (MTD).....	6
Jackson.....	7	Glorion.....	5½	Queen Turf.....	6
Sears.....	7	Scott.....	5½	Cyclone.....	5½

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 12¾ oz. per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

FOR ESTABLISHED TURF

With a calibrated spreader, apply evenly at rate of 8 lbs. per 1000 sq. ft. Treated areas should be lightly watered after application if no rain falls within one week.

ANNUAL BLUEGRASS (POA ANNUA) CONTROL

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turf areas. Irrigation will encourage the existing turfgrasses to fill in these thin areas formerly occupied by annual bluegrass. Apply 12 lbs. per 1000 sq. ft. (1½ oz. per 100 sq. ft.) in late summer or early fall (mid July to mid September) or in early spring prior to annual bluegrass germination.

CAUTION

May be harmful if swallowed. Avoid contact with skin. Avoid contamination of food-stuffs, feeds, and fishponds. Wash thoroughly with soap and water after using.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Avoid prolonged or repeated inhalation of dust.

Do not apply to dichondra turf. Do not apply in spring to turfgrass planted the previous fall.

Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

WARRANTY

Lebanon Chemical Corporation, the manufacturer, warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.


TREATS 5,000 SQ. FT.

LEBANON

ACCEPTED

Jan. 28, 1978

UNDER THE PROVISIONS OF THE
FEDERAL FOOD, DRUG AND COSMETIC ACT
FOR THE CONTROL OF MISBRANDED
DRUGS, THIS PRODUCT IS SUBJECT
TO THE FOLLOWING COMMENTS:
961-248



WITH
CRABGRASS
PREVENTION



10-0
WI
CRABGRASS
Contains

A pre-emergence herbicide applied
crabgrass, poa annua, goosegrass, watergrass

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-a,a,a-trifluoro-2,6-

WITH CRABGRASS CONTROL

10-6-4

10-6-4 WITH CRABGRASS CONTROL Contains BALAN™

A pre-emergence herbicide applied directly to established turf for control of crabgrass, poa annua, goosegrass, watergrass, yellow and green foxtail.

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine 0.43%

INERT INGREDIENTS: 99.57%

(Relative to pesticide content) TOTAL 100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN (See back panel for additional cautions)

BALAN™ - the trade mark for BENEFIN-ELANCO PRODUCTS COMPANY
A DIVISION OF ELI LILLY & COMPANY

E.P.A. Reg. No. 961-248

ELANCO CHEMICAL CORPORATION LEBANON, PA. 17042

LEBANON GREEN GOLD 10-6-4

WITH CRABGRASS CONTROL

GUARANTEED MINIMUM FERTILIZER ANALYSIS

TOTAL NITROGEN (N)	10.0%
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	6.0%
SOLUBLE POTASH (K ₂ O)	4.0%
Potential acidity 600 lb. calcium carbonate per ton.	

LEBANON CHEMICAL CORPORATION

LEBANON, PA. 17042

Feeds your lawn, kills crabgrass upon germination. Grass seed can be applied 60 days after applying GREEN GOLD WITH CRABGRASS CONTROL.

ESTABLISHES A WEED CONTROL ZONE

GREEN GOLD WITH CRABGRASS CONTROL establishes a weed control zone by killing certain annual weed seeds as they germinate. Application prior to weed germination will prevent poa annua, crabgrass, goosegrass, watergrass, yellow and green foxtail.

DIRECTIONS FOR USE

GREEN GOLD WITH CRABGRASS CONTROL can be used on a variety of turf or mixture of turf grasses.

Bluegrasses, ryegrass, fescue, bermudagrass, zoysiagrass, St. Augustinegrass, centipede-grass and bahiagrass are tolerant. Do not apply to newly sprigged areas of St. Augustine, Bermuda, or Centipede grasses.

SPREADER SETTINGS

Lawn Beauty.....	7	Central.....	5½	Modern (MTD)	6
Jackson.....	7	Glorion	5½	Queen Turf	6
Sears	7	Scott	5½	Cyclone	5½

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 12¾ oz. per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

FOR ESTABLISHED TURF

With a calibrated spreader, apply evenly at rate of 8 lbs. per 1000 sq. ft. Treated areas should be lightly watered after application if no rain falls within one week.

ANNUAL BLUEGRASS (POA ANNUA) CONTROL

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turf areas. Irrigation will encourage the existing turfgrasses to fill in these thin areas formerly occupied by annual bluegrass. Apply 12 lbs. per 1000 sq. ft. (19 oz. per 100 sq. ft.) in late summer or early fall (mid July to mid September) or in early spring prior to annual bluegrass germination.

CAUTION

May be harmful if swallowed. Avoid contact with skin. Avoid contamination of food-stuffs, feeds, and fishponds. Wash thoroughly with soap and water after using.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Avoid prolonged or repeated inhalation of dust.

Do not apply to dichondra turf. Do not apply in spring to turfgrass planted the previous fall.

Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

WARRANTY

AC
Jan 28, 1972
961-248

GREEN GOLD 10-6-4

AC

WITH CRABGRASS CONTROL

Jan 28, 1977

GUARANTEED MINIMUM FERTILIZER ANALYSIS

TOTAL NITROGEN (N)	10.0%
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	6.0%
SOLUBLE POTASH (K ₂ O)	4.0%
Potential acidity 600 lb. calcium carbonate per ton.	

961-248

LEBANON CHEMICAL CORPORATION

LEBANON, PA. 17042

Feeds your lawn, kills crabgrass upon germination. Grass seed germinates 10 to 14 days after applying GREEN GOLD WITH CRABGRASS CONTROL.

ESTABLISHES A WEED CONTROL ZONE

GREEN GOLD WITH CRABGRASS CONTROL establishes a weed control zone by killing certain annual weed seeds as they germinate. Application prior to weed germination will prevent poa annua, crabgrass, goosegrass, watergrass, yellow and green foxtail.

DIRECTIONS FOR USE

GREEN GOLD WITH CRABGRASS CONTROL can be used on a variety of turf or mixture of turf grasses.

Bluegrasses, ryegrass, fescue, bermudagrass, zoysiagrass, St. Augustinegrass, centipede-grass and bahiagrass are tolerant. Do not apply to newly sprigged areas of St. Augustine, Bermuda, or Centipede grasses.

SPREADER SETTINGS

Lawn Beauty.....	7	Central.....	5½	Modern (MTD)	6
Jackson.....	7	Glorion	5½	Queen Turf	6
Sears	7	Scott	5½	Cyclone	5½

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 12¾ oz. per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

FOR ESTABLISHED TURF

With a calibrated spreader, apply evenly at rate of 8 lbs. per 1000 sq. ft. Treated areas should be lightly watered after application if no rain falls within one week.

ANNUAL BLUEGRASS (POA ANNUA) CONTROL

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turf areas. Irrigation will encourage the existing turfgrasses to fill in these thin areas formerly occupied by annual bluegrass. Apply 12 lbs. per 1000 sq. ft. (19 oz. per 100 sq. ft.) in late summer or early fall (mid July to mid September) or in early spring prior to annual bluegrass germination.

CAUTION

May be harmful if swallowed. Avoid contact with skin. Avoid contamination of food-stuffs, feeds, and fishponds. Wash thoroughly with soap and water after using.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Avoid prolonged or repeated inhalation of dust.

Do not apply to dichondra turf. Do not apply in spring to turfgrass planted the previous fall.

Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

WARRANTY

Lebanon Chemical Corporation, the manufacturer, warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.