

Net Weight: 25 Lbs.

Treats: 25,000 Sq. Ft.

CHINCH BUG
~~INSECT AND GRUB~~
GREENSKEEPER CONTROL
with Dursban*

CONTROLS

Chinch Bugs, Sod Webworms (Lawn Moths), Ants, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Sowbugs (Pillbugs), Fleas, Chiggers, Clover Mites, Grasshoppers, Armyworms and Hyperodes (Turfgrass) Weevils infesting lawns and other ornamental areas. Controls larvae of Japanese Beetle and European Chafer.

ACTIVE INGREDIENT:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 2.32%

INERT INGREDIENTS:

97.68%

TOTAL: 100.00%

Dursban® - a Trademark of DowElanco	EPA Reg. No. 961-261	(L) (D) EPA Est. 961-PA-01:961-IL-01
U.S. Patent No. 3,244,586		Superscript used is first letter of lot number.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN
WARNING

May be harmful if swallowed. Do not take internally. Avoid inhaling dust. Avoid eye contact. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control center. Drink one or two glasses of water, and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air if symptoms of cholinesterase inhibition appear, and get medical attention immediately.

IF ON SKIN: Wash thoroughly with soap and water. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection, is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

A C C E P T E D
APR 18 1995
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 961-261

BEST COPY AVAILABLE

1982

Apply this product uniformly over the turf area at a rate of 1 lb. per 1,000 sq. ft. Make the first treatment when insects first appear and re-treat as needed. Keep children and pets off treated area until material is washed into the soil with 30 gallons of water per 1,000 sq. ft. and grass is dry.

To control Hyperodes (Turfgrass) Weevils, use twice the rate of this product indicated for other insects. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for the best time to treat in your area.

For control of larvae of Japanese Beetle and European Chafer, apply at the rate of 2 to 4 lbs. per 1,000 sq. ft. Treat when white grubs are young and feeding near the soil surface, usually during late July and August, or as recommended by your local Agricultural Extension Service Specialist.

Immediately after application for weevil or larvae, wash granules into the soil with 1/2 to 1 inch of water.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

SPREADER SETTINGS

	<u>1 Lb.</u>	<u>2 Lb.</u>
Show 'N Tell	6	7
Ortho Rotary	3	5
Scott	5	6
Earthway Rotary	14	16
Scott Rotary	E	H
Glorion	5	6
Sears	5 1/2	6 1/2
Central	5 1/2	6
MTD	2	2 1/2
Jackson	5 1/2	6
Jet Spred®	3	4
Lesco	F	H
Ortho Drop	5	7
Cyclone	3	4

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply only 16 oz. per 100 sq. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

BEST COPY AVAILABLE