

TREATS 5,000 SQ. FT.

NET WEIGHT 20 LBS.

LEBANON WEED AND FEED 10-6-4 WITH 2,4-D AND MCPP

MINIMUM GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	10.0%
Available Phosphoric Acid (P ₂ O ₅)	6.0%
Soluble Potash (K ₂ O)	4.0%
Potential Acidity 1,000 lbs. Calcium Carbonate per ton.	

LEBANON CHEMICAL CORPORATION
LEBANON, PA. 17042

FOR A LUXURIOUS, WEED FREE LAWN

The 10-6-4 fertilizer stimulates grass growth to quickly conceal spots where weeds have died. Kills weeds and grows healthy green grass in one application.

EFFECTIVE WEED CONTROL

Kills many common weeds, including buckhorn, dandelion, chickweed, clover, ground ivy, knock-weed, plantain. knot-

SAFE FOR ESTABLISHED LAWNS

Will not harm most lawn grasses. May cause some temporary browning, particularly on Seaside, Arlington and Congressional bent grasses. If browning is noted, water thoroughly to expedite recovery.

SPREADER SETTINGS


Lawn Beauty	5	Central	6	Sears	5
MTD	5	Queen Turf	5	Jackson	6
Cyclone	4	Scott	4 1/2	Glorion	6

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 6 oz. per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the spreader settings necessary to apply the correct amount.

When applying with a spinner type spreader, use extreme caution to avoid any contact or application to desirable plants.

DIRECTIONS

The best time to destroy weeds is when they are young and actively growing, before they crowd out desirable grasses. Apply at the rate of 4 lbs. per 1,000 sq. ft., in Spring or early Fall, when ground moisture is adequate and weeds are actively growing. If used during dry period, water lawn thoroughly 24 hours prior to application.



LEBANON, PA. 17042

FOR A LUXURIOUS, WEED FREE LAWN

The 10-6-4 fertilizer stimulates grass growth to quickly conceal spots where weeds have died. Kills weeds and grows healthy green grass in one application.

EFFECTIVE WEED CONTROL

Kills many common weeds, including blackberry, dandelion, chickweed, clover, ground ivy, knockweed, plantain.

knock-

SAFE FOR ESTABLISHED LAWNS

Will not harm most lawn grasses. May cause some temporary browning, particularly on Seaside, Arlington and Congressional bent grasses. If browning is noted, water thoroughly to expedite recovery.

SPREADER SETTINGS

Lawn Beauty	5	Central	6	Stans	5
MTD	5	Queen Turf	5	Jackson	6
Cyclone	4	Scott	4 1/2	Glorion	6

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 6 oz. per 100 sq. ft. Rate of soil, moisture and spreader condition can vary the spreader settings necessary to apply the correct amount.

When applying with a spinner type spreader, use extreme caution to avoid any contact or application to desirable plants.

DIRECTIONS

The best time to destroy weeds is when they are young and actively growing, before they crowd out desirable grasses. Apply at the rate of 4 lbs. per 1,000 sq. ft. in Spring or early Fall, when ground moisture is adequate and weeds are actively growing. If used during dry period, water lawn thoroughly 24 hours prior to application.

For persistent weeds, when weeds reappear, or if a heavy rain occurs within 2 days after application, a second application may be necessary.

Apply to new lawns only after 2nd or 3rd mowing. Do not apply to very young grass, or to predominantly bent grass turf.

Do not mow, water, or make heavy use of lawn for at least 24 hours after application.

CAUTION

Avoid inhaling dust. Avoid contact with skin, eyes, or clothing. In case of eye contact, flush with water. Vapors from this product may injure susceptible plants in the immediate vicinity. Bentgrass, St. Augustine grass, Dichondra, Clover and Carpetgrass may be injured. Avoid drift to susceptible plants, as this product may injure flowers, ornamentals, and vegetables.

Do not store near fertilizers, seeds, insecticides or fungicides. Clean spreader thoroughly after use.

Do not reuse container. Destroy when empty. Bury in a safe place. Do not burn.

Do not contaminate water by cleaning of equipment or disposal of wastes.

WARRANTY

Lebanon Chemical Corporation warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

RECOMMENDED

10-

BUCK
KNO
CAUT

erary browsing, particularly on Swards,
is noted, water thoroughly to expedite

INGS

Stans _____ 5
Jackson _____ 6
Glorton _____ 6

in 100 sq. ft. and adjust to apply 6 oz. per 100
water settings necessary to apply the correct

any contact or application to desirable plants.

NS

actively growing, before they crowd out
in Spring or early Fall, when ground
used during dry period, water lawn

rain occurs within 2 days after applica-

ply to very young grass, or to predomi-

24 hours after application.

ic. In case of eye contact, flush with water.

in the immediate vicinity. Benzene, St.
injured. Avoid drift to susceptible plants,
stables.

dos. Clean spreader thoroughly after use.

in place. Do not burn.

disposal of wastes.

to its chemical description and is reasonably fit
actions under normal conditions of use, but neither
particular purpose, express or implied, extends
normal conditions, or under conditions not reason-

**D
AND
F
M
D
I
O
O
N**

FEED

10-6-4 WITH 2,4-D AND MCPP

CONTROLS

**BUCKHORN, DANDELION, CHICKWEED, CLOVER, GROUND IVY,
KNOTWEED, PLANTAIN, AND OTHER BROADLEAF WEEDS**

CAUTION: KEEP OUT OF REACH OF CHILDREN (SEE BACK PANEL FOR ADDITIONAL CAUTIONS)

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid* 0.858%
Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid** .. 0.868%

INERT INGREDIENTS:

..... 98.264%

TOTAL 100.000%

*Equivalent to 0.72% 2,4-dichlorophenoxyacetic acid

**Equivalent to 0.72% 2-(2-methyl-4-chlorophenoxy) propionic acid

E.P.A. Reg. No. 361-263

LEEBYANON CHEMICAL CORPORATION

ACCEPTED

18 1972

**INSECTICIDE
HERBICIDE AOT**