



NET WT. 10 LBS.



96/273

PREEMERGENCE WEED CONTROL

LAWNS-SHRUBBERY-VEGETABLES-FLOWERS

Controls Crabgrass, Poa Annua (Annual Bluegrass), Carpetweed, Common Chickweed, Lambsquarters, Purslane, Green and Yellow Foxtail, Lovegrass, and Witchgrass.



CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions

ACTIVE INGREDIENT:

PREEMERGENCE WEED CONTROL

LAWNS-SHRUBBERY-VEGETABLES-FLOWERS

Controls Crabgrass, Poa Annua (Annual Bluegrass), Carpetgrass, Common Chickweed, Lambsquarters, Purslane, Green and Yellow Foxtail, Lovegrass, and Witchgrass.



CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions

ACTIVE INGREDIENT:		
Dimethyl ester of tetrachloroterephthalic acid*	5.0%	
INERT INGREDIENT:	95.0%	
	TOTAL	100.0%

E.P.A. REG. NO. 961-273

*Use as a pre-emergence herbicide patented under U.S. Patent No. 2,923,634

LEBANON CHEMICAL CORP.
LEBANON, PA. 17042

PREEMERGENCE WEED CONTROL

Up to 4 months of weed control may be expected from the use of LEBANON PREEMERGENCE WEED CONTROL at recommended rates. For cultivated areas, apply to weed-free surface within two to three days following cultivation to prevent weed seed germination. Disturbing soil surface following application will reduce effectiveness. If no rain occurs within three or four days after application, it is advisable to sprinkle area, applying at least 1/2 inch of water.

WEEDS CONTROLLED

Crabgrass, green and yellow foxtail, carpetweed, common chickweed, lambsquarters, lovegrass, purslane, witchgrass, barnyard grass, goosegrass, nodding spurge, redroot pigweed.
Annual bluegrass.

DIRECTIONS

LAWNS

Spring Application: 10 lbs. cover 2100 sq. ft. Apply LEBANON PREEMERGENCE WEED CONTROL at rate of 4 1/2 lbs. per 1000 sq. ft. early in spring, before crabgrass and other annual grasses and weeds germinate. One application, before Pernettya becomes tall, usually provides control for the entire season. If application is made on a new lawn, grass seed-bags should be at least two inches in height and uniformly green. Early spring application allows fall sowing of permanent grass. Delay seeding for at least 60 days after application of LEBANON PREEMERGENCE WEED CONTROL.

CAUTION: Do not apply to Dickcissel lawns. Do not apply to greens planted in Colansey or Toronto berms. Use no more than the recommended rate.

LATE SUMMER OR EARLY FALL APPLICATION: 10 lbs. cover 1400 sq. ft. To control annual bluegrass and other late-germinating grasses and weeds, an application in late summer or early fall of 7 lbs. per 1000 sq. ft. should follow the spring application. Consult agricultural department station or extension service weed specialists for proper application time in your area.

Suggested spreader settings for application rate of 4 1/2 lbs. per 1000 sq. ft. are listed below. These are only approximate settings, and are not intended to relate to all spreaders. Check your own spreader on a small area by using an appropriate amount. Rate of walk, condition of spreader, and other factors influence the rate of flow.

Levon Heavy	7	Autumn	8	Gleason	8
MTD	8	Scott	7	Bears	8
Queen Turf	8	Control	8	Cyclone	4.5

FLOWER AND ORNAMENT BEDS: 10 lbs. cover 2200 sq. ft.

Apply at the rate of 4 1/2 lbs. per 1000 sq. ft. Apply immediately after establishment and planting of bed. Weed control for up to 4 months may be expected if properly applied to weed-free cultivated bed. Plants must be well-established before application.

LEBANON PREEMERGENCE WEED CONTROL is specifically recommended for weed control on the following:

Abelia	Columbine	Forsythia	Ivy	Moss rose	Snake's head
Aster	Delphinium	Geranium	Japanese maple	Mother-of-pearl	Spiraea
Asplenium	Drumstick	Hydrangea	Japanese spirea	Thyme	Strawflower
Azalea	Flax	Lily	Japanese anemone	Mountain laurel	Sunflower
Begonia	Flowering quince	Lily	Japanese iris	Mourning glories	Sweet pea
Begonia	Flowering quince	Lily	Japanese iris	Nasturtium	Sycamore
Begonia	Flowering quince	Lily	Japanese iris	Pachysandra	Tree peony
Begonia	Flowering quince	Lily	Japanese iris	Pony	Tulip tree
Begonia	Flowering quince	Lily	Japanese iris	Persea	Verbena
Begonia	Flowering quince	Lily	Japanese iris	Pink dogwood	Viburnum
Begonia	Flowering quince	Lily	Japanese iris	Rhododendron	Violet
Begonia	Flowering quince	Lily	Japanese iris	Rose	Weigelia
Begonia	Flowering quince	Lily	Japanese iris	Russian olive	Yarrow fern
Begonia	Flowering quince	Lily	Japanese iris	Sage, scarlet	Yew
Begonia	Flowering quince	Lily	Japanese iris	Salvia	Zinnia

Do not use on the following plants:

Begonia	Carnation	Geranium	Pansy	Sweet William
Butter plant	Osiris	Meibomia-anthemum	Phlox	Telanthra

VEGETABLE GARDENS: The following crops may be treated with LEBANON PREEMERGENCE WEED CONTROL as described.

BROCCOLI, BRUSSELS SPROUTS, CALLIFLOWER, CABBAGE: Apply 4 to 4 1/2 lbs. per 1000 sq. ft. uniformly to the soil at seeding or transplanting. LEBANON PREEMERGENCE WEED CONTROL can be applied directly over transplants without injury. If weeds have emerged, crops should be cleanly cultivated before application.

BEANS (DRY OR SNAP) AND SOUTHERN PEAS (BLACKEYED): Apply uniformly to the soil at seeding at the rate of 4 to 4 1/2 lbs. per 1000 sq. ft.

BEETED CHIMBARRS, SQUASH (WINTER AND SUMMER), MELONS (CANTALOUPE, HONEYDEW, WATERMELON): Apply 4 to 4 1/2 lbs. per 1000 sq. ft. uniformly to the soil 2 to 6 weeks after seeding. Cultivate the crops before application. If weeds have emerged.

COLLARDS, KALE, MUSTARD GREENS, TURNIPS (GREEN AND ROOTS): Apply uniformly to the soil at the rate of 4 to 4 1/2 lbs. per 1000 sq. ft. at seeding.

CALCIC AND ONIONS: Apply 4 to 4 1/2 lbs. per 1000 sq. ft. uniformly to the soil at seeding, transplanting or in rows at lay by. LEBANON PREEMERGENCE WEED CONTROL can be applied directly over transplants without injury. Lay by applications can be made on onions up to 14 weeks after seeding or transplanting. If weeds emerge prior to lay by, cultivate the crop before application.

ROTAMIS (WHOLE OR CUT PIECES): Apply uniformly to the soil at 4 to 4 1/2 lbs. per 1000 sq. ft. during planting, drop off or lay by. Lay by application can be made up to nine weeks after planting. If weeds have emerged at lay by, the crop should be cultivated before application. The initial sprayer irrigation should not exceed one inch. LEBANON PREEMERGENCE WEED CONTROL is not recommended for use on potatoes in California.

SWEET POTATOES AND YAMS: Apply 4 to 4 1/2 lbs. per 1000 sq. ft. uniformly to the soil at transplanting or lay by. LEBANON PREEMERGENCE WEED CONTROL can be applied directly over transplants without injury. Lay by applications can be made up to six weeks after transplanting. If weeds have emerged, cultivate the crop before applying LEBANON PREEMERGENCE WEED CONTROL.

TOMATOES, EGGPLANT, PEPPERS: Uniform application to the soil of 4 to 4 1/2 lbs. per 1000 sq. ft. should be effected to a depth of four to six weeks after transplanting. Plants should be well established and growing in conditions favorable to LEBANON PREEMERGENCE WEED CONTROL can be applied directly over the plants without injuring them. Cultivate or weed the crop before treatment if weeds have emerged.

STRAWBERRIES: New plantings — Apply 4 lbs. per 1000 sq. ft. uniformly to soil at transplanting. Established plantings — Apply in early fall and early spring at the rate of 4 lbs. per 1000 sq. ft. to control fall and early spring germinating weeds. Do not apply after first bloom. These applications can be made directly over plants without injury.

SOYBEANS: Apply 4 to 4 1/2 lbs. per 1000 sq. ft. uniformly to the soil at seeding.

REPLANTING

Replanting crops other than those included on this label in LEBANON PREEMERGENCE WEED CONTROL may result in crop injury within 6 months of application. If replanting is required due to early crop failure, the planting of onions, seeded melons, potatoes, tomatoes, eggplant or peppers at this time may result in crop injury. If weeds are present on the label may be planted following harvest of a LEBANON PREEMERGENCE WEED CONTROL treated crop.

CAUTION

Keep out of reach of children.

Harmful if swallowed.

Avoid contact with eyes and skin. In case of contact, flush eyes with plenty of water. Wash after handling or using.

Do not exceed recommended application rates.

Do not store near food and foodstuffs, seed, fertilizers or other pesticides in order to prevent contamination.

See cautions under laws recommendations.

Store in dry place.

Do not graze on treated areas or feed plant refuse to livestock.

Do not reuse container. Destroy when empty.

WARRANTY

Manufacturer warrants that this product, if used in accordance with the directions on the label, will provide control of the weeds listed on the label. If the weeds listed on the label are not controlled, the manufacturer will refund the purchase price of the product. This warranty does not apply if the product is used in a manner not recommended on the label, or if the product is used on a crop not listed on the label. The manufacturer does not warrant that the product will control weeds not listed on the label.