

PM 19 961-275

10/11

NET WEIGHT: 20 LBS. (9.08 Kg)

TREATS: 5,000 Sq. Ft.

# GREEN GOLD LAWN FOOD

# 19-4-4<sub>with</sub>

# INSECT AND GRUB CONTROL

Contains Dursban\*

Controls: Chinch Bugs, Sod Webworms (Lawn Moths), Ants, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Sowbugs (Pillbugs), Fleas, Chiggers, Clover Mites, Grasshoppers, Armyworms and Hyperodes (Turfgrass) Weevils infesting home lawns and other ornamental turf areas. Also controls grubs (larvae) of Japanese Beetles and European Chafer.

**ACTIVE INGREDIENT**

Chlorpyrifos[0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl)phosphorothioate]\* 0.60%

**INERT INGREDIENTS** 99.40%

**TOTAL:** 100.00%

\*Dursban - a trademark of DowElanco  
U.S. Patent No. 3,244,586

EPA Reg. No. 961-275

(L) (D)  
EPA Est. 961-PA-01:961-IL-01  
Superscript used is first letter of lot number.

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
See Back Panel for Additional Precautionary Statements

**ACCEPTED**  
**APR 18 1995**  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 961-275

**MANUFACTURED BY**  
**Lebanon Chemical Corporation**  
**Lebanon, PA 17042**

Page 1 of 4 Pages  
961-275  
01/10/95

**BEST COPY AVAILABLE**

134

Scj

### GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	19.0%
3% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	4.0%
Soluble Potash (K <sub>2</sub> O)	4.0%

PRIMARY NUTRIENTS FROM: Urea, Methylene Ureas, Ammonium Phosphate, Ammonium Sulfate, Muriate of Potash.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

**KEEP OUT OF REACH OF CHILDREN.** Harmful if swallowed. Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before re-use. Keep away from food, feedstuffs and domestic water supplies. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

**NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection, is an antidote.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Induce vomiting immediately by giving two glasses of water and sticking finger down throat. Call a physician.

**IF INHALED:** Remove victim to fresh air. If symptoms occur, get medical attention.

**IF ON SKIN:** Wash thoroughly with soap and water.

**IF IN EYES:** Flush with plenty of water for 15 minutes and get medical attention.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

3/4

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.**

## APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL

Feeds your lawn the essential food elements required for good growth while controlling the following lawn and home pests: Chinch Bugs, Sod Webworms (Lawn Moths), Ants, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Sowbugs (Pillbugs), Fleas, Chiggers, Clover Mites, Grasshoppers, Armyworms, and Hyperodes (Turfgrass) Weevils. Also for control of larvae of Japanese Beetle and European Chafer.

For control of all insects named above, except Hyperodes Weevils and larvae of Japanese Beetles and European Chafer, apply at the rate of 4 lbs. per 1,000 sq. ft. to turf or lawn area. Treat when insects first appear, re-treat as needed.

For control of Hyperodes (Turfgrass) Weevils, make a double application using 8 lbs. per 1,000 sq. ft. Treat suspected problem areas in mid-April and again in mid-May. Consult your Agricultural Extension Service specialist for best time to treat in your area.

For control of larvae of Japanese Beetle and European Chafer, apply at the rate of 8 lbs. per 1,000 sq. ft. (use 2 successive applications per table directions). Treat when white grubs are young and actively feeding near the soil surface, usually during late July and August, or as recommended by your local Agricultural Extension Service Specialist.

Immediately after all applications, wash granules into the soil with 1/2 to 1 inch of water.

## STORAGE AND DISPOSAL

**STORAGE:** Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

Page 3 of 4 Pages

981-275

01/10/95

**BEST COPY AVAILABLE**

3

42/4

## WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

Page 4 of 4 Pages  
961-275  
01/10/95

BEST COPY AVAILABLE

4