

Net Weight 25 lbs.

Covers 5,000 sq. ft.

LEBANON LAWN INSECT AND GRUB CONTROL

Contains 2% Diazinon® insecticide

WHEN USED AS DIRECTED, KILLS the following insects:

- | | | | |
|-----------------|--------------|------------------------------|-----------------|
| Ants | Clover mites | Lawn moths
(Sod webworms) | Sowbugs |
| Armyworms | Crickets | Leafhoppers | Ticks |
| Billbugs | Cutworms | Millipedes | White grubs |
| Brown dog ticks | Digger wasps | Springtails
(Collembola) | Japanese beetle |
| Chiggers | Earwigs | | European chafer |
| Cinch bugs | Fleas | | Southern chafer |


*Diazinon is trademark of CIBA-GEIGY Corp.

ACTIVE INGREDIENT:
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate*

INERT INGREDIENTS:

TOTAL

E.P.A. Reg. No. 961-276



CAUTION: KEEP OUT OF REACH OF CHILDREN. For additional cautions see below.


DIRECTIONS

Application of **LEBANON LAWN INSECT AND GRUB CONTROL** at the rate of 5 lbs. per 1,000 sq.ft. will control the following insects: Ants, Armyworms, Brown dog ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (Sod webworms), Leafhoppers, Millipedes, Springtails (Collembola), Sowbugs and Ticks. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary.

To control Billbugs and White grubs of Japanese beetle, European chafer, Southern chafer, apply **LEBANON LAWN INSECT AND GRUB CONTROL** at the rate of 6 lbs. per 1,000 sq. ft. Water grass thoroughly after application. For grub control, make application any time between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

For Hyperodes Weevil (New York only): Apply 5 lbs. per 1,000 sq.ft. Water grass thoroughly after application. Treat problem area in mid-April and again in mid-May.

Application is best made with a fertilizer spreader. Be sure to apply at the recommended rate regardless of method of distribution. The settings given below are approximate, and may require variation due to the age of your spreader, the rate of walk, and other possible factors. Always calibrate your own spreader on a small area (100 sq. ft) to apply the right amount.



Spreader	5 lbs./1,000 sq. ft.	6 lbs./1,000 sq. ft.
Jackson	6	6½
Sears	6	6½
Scott	5	6
Lawn Beauty	6	6½
Central	5	6
MTD	5	6
Cyclone	5	5½
Glorion	5	6

SPOT TREATMENT

Ants: Apply one tablespoon of **LEBANON LAWN INSECT AND GRUB CONTROL** over and around each individual ant hill.

Digger wasps: Apply one tablespoon of **LEBANON LAWN INSECT AND GRUB CONTROL** in and around each individual ground nest opening. Apply around dusk when wasps are not active.

FOUNDATION BAND TREATMENT

The following insects are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.

- | | |
|-----------------|--------------|
| Ants | Fleas |
| Brown dog ticks | Millipedes |
| Clover mites | Sowbugs |
| Crickets | Springtails |
| Earwigs | (Collembola) |

To control the above insects, apply a five-foot band of **LEBANON LAWN INSECT AND GRUB CONTROL** next to the foundation of the house. Distribute the granules uniformly at the rate of 8 ounces for each 5 foot by 20 foot area.

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes and clothing. May be absorbed through the skin. Wash thoroughly after handling.

Cinch bugs

Fleas

(Collembola)

Southern chafer

*Miazinon is trademark of CIBA-GEIGY Corp.

ACTIVE INGREDIENT:
 O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate*
INERT INGREDIENTS:
TOTAL

ACCEPTED
 # 961-276
 2.0%
 98.8%
 100%
 MAY 29 1973
 UNITED STATES FEDERAL INSECTICIDE
 AND RODENT CONTROL ACT
 POISON REGISTER
 SUBJECT

E.P.A. Reg. No. 961-276

CAUTION: KEEP OUT OF REACH OF CHILDREN. For additional cautions see below.

DIRECTIONS

Application of LEBANON LAWN INSECT AND GRUB CONTROL at the rate of 5 lbs. per 1,000 sq.ft. will control the following insects: Ants, Armyworms, Brown dog ticks, Chiggers, Cinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (Sod webworms), Leafhoppers, Millipedes, Springtails (Collembola), Sowbugs and Ticks. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary.

To control Billbugs and White grubs of Japanese beetle, European chafer, Southern chafer apply LEBANON LAWN INSECT AND GRUB CONTROL at the rate of 6 lbs. per 1,000 sq. ft. Water grass thoroughly after application. For grub control, make application any time between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

For Hyperodes Weevil (New York only): Apply 5 lbs. per 1,000 sq.ft. Water grass thoroughly after application. Treat problem area in mid-April and again in mid-May.

Application is best made with a fertilizer spreader. Be sure to apply at the recommended rate regardless of method of distribution. The settings given below are approximate, and may require variation due to the age of your spreader, the rate of walk, and other possible factors. Always calibrate your own spreader on a small area (100 sq. ft) to apply the right amount.

ACCEPTED
 MAY 29 1973
 UNITED STATES FEDERAL INSECTICIDE
 AND RODENT CONTROL ACT
 POISON REGISTER
 SUBJECT

Spreader	5 lbs./1,000 sq. ft.	6 lbs./1,000 sq. ft.
Jackson	6	6 1/2
Sears	6	6 1/2
Scott	5	6
Lawn Beauty	6	6 1/2
Central	5	6
MTD	5	6
Cyclone	5	5 1/2
Glorion	5	6

SPOT TREATMENT
Ants: Apply one tablespoon of LEBANON LAWN INSECT AND GRUB CONTROL over and around each individual ant hill.

Digger wasps: Apply one tablespoon of LEBANON LAWN INSECT AND GRUB CONTROL in and around each individual ground nest opening. Apply around dusk when wasps are not active.

FOUNDATION BAND TREATMENT

The following insects are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.

- | | |
|-----------------|--------------|
| Ants | Fleas |
| Brown dog ticks | Millipedes |
| Clover mites | Sowbugs |
| Crickets | Springtails |
| Earwigs | (Collembola) |

To control the above insects, apply a five-foot band of LEBANON LAWN INSECT AND GRUB CONTROL next to the foundation of the house. Distribute the granules uniformly at the rate of 8 ounces for each 5 foot by 20 foot area.

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes and clothing. May be absorbed through the skin. Wash thoroughly after handling.

Avoid contamination of feed and foodstuffs.

Treat lawns in late afternoon. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

Do not re-use this container. Destroy and bury, or discard in a safe place.

WARRANTY

Lebanon Chemical Corporation warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and buyer assumes the risk of any such use.

LEBANON CHEMICAL CORPORATION

LEBANON, PA. 17042