

# LEBANON TURF FUNGICIDE

NET WEIGHT 10 LBS.

Treats 2,500 sq. ft. at Preventative Rate

CONTAINS DACONIL 2787 fungicide, an effective broad spectrum fungicide for the control of many turf diseases, including the following:

HELMINTHOSPORIUM LEAF SPOT, MELTING OUT, CURVULARIA (Copper Spot), CUPPER SPOT, Gaeumannomyces (Dollar Spot), Sclerotinia (Brown Patch), Rhizoctonia (Stem Rust), GRAM LEAF SPOT, and Anthracnose and STEM RUST of bluegrass.

### SPREADER SETTINGS

The following spreader settings are approximate and should be used only as a guide in setting your own spreader to deliver at the rate of 4 pounds per 1,000 sq. ft. Many factors affect the rate of individual spreaders, such as rate of walk, age and condition of spreader, etc.

Green	6	Green	7 1/2	Aspen	7 1/2
Cobra	7 1/2	Wick	7 1/2	Sears	7 1/2
Scor	8 1/2	Wick	7	Cruise	4 1/2

### DIRECTIONS

Apply when grass is damp, as after a rain or in the early morning. Following application, do not water or mow grass for a minimum of 24 hours. Use 4 pounds per 1,000 sq. ft. for normal control. Increase rate to 6 pounds per 1,000 sq. ft. for severely affected areas.

For control of Helminthosporium Leaf Spot, Melting Out and Curvularia, apply in early Spring. Continue applications at 7 to 10 day intervals during wet season.

To control Copper Spot, Dollar Spot, Brown Patch, Gray Leaf Spot and Stem Rust, apply in early Spring to early fall. Continue applications at 7 to 14 day intervals into early fall.


### CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using.

This product is toxic to fish. Keep out of lakes, streams, or ponds.

TREATS 2,500 SQ. FT.  
PREVENTATIVE RATE

NET WT. 10 LBS.



TURF FUNGICIDE

# TURF FUNGICIDE

TURF FUNGICIDE

CONTAINS  
**\*DACONIL 2787\***

normal control, increase rate to 6 pounds per 1,000 sq. ft. for severely affected areas.

For control of Helminthosporium Leaf Spot, Melting Out, and Curvularia, apply in early Spring. Continue applications at 7 to 10 day intervals during wet season.

To control Copper Spot, Dollar Spot, Brown Patch, Gray Leaf Spot, and Stem Rust, apply in early Spring to early Summer, and continue applications at 7 to 14 day intervals into early Fall.

### CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not pollute surface water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not graze treated areas or feed clippings to livestock.

Do not reuse empty containers. Destroy by burying with waste material or by other means to prevent reuse.

### WARRANTY

This product is warranted to perform as described on the label when used in accordance with the directions on the label. The user assumes all responsibility for the results obtained. The manufacturer does not warrant the results of use.

ACCEPTED

March 11, 1974

961-277

FUNGICIDE

CONTAINS  
**\*DACONIL 2787\***

TURF

LEBANON

Used as directed, will prevent turf problems caused by Curvularia, Leaf Spot Melting Out, Copper Spot, Dollar Spot, Brown Patch, Gray Leaf Spot, and Stem Rust of bluegrass.

**CAUTION:** Keep out of reach of children. See back panel for additional cautions.

#### ACTIVE INGREDIENTS:

Chlorothalonil (Tetrachloroisophthalonitrile) 50%

#### INERT INGREDIENTS

TOTAL 100%

EPA Reg. No. 961-277  
EPA Est. 961-PA-1

\*Covered under U.S. Patent No. 3,290,355

Formulation of Chlorothalonil

MANUFACTURED BY  
**LEBANON CHEMICAL CORPORATION**  
**LEBANON, PA. 17042**