

GREENVIEW
EVER PREEN

15/91
- 3
961-283

WEED CONTROL PLUS FERTILIZER
FOR EVERGREENS-SHRUBS-FLOWERS-VEGETABLES

ACTIVE INGREDIENT:

Trifluralin (a,a,a-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine) 0.74%

INERT INGREDIENTS 99.26%

EPA Reg. No. 961-283

ACCEPTED
OCT 29 1986
Use on the Federal Insecticide, Fungicide, and Rodenticide Act as a Fertilizer
registered under
EPA Reg. No. 961-283

GUARANTEED FERTILIZER ANALYSIS


Total Nitrogen (N)	%
Available Phosphoric Acid (P ₂ O ₅)	%
Soluble Potash (K ₂ O)	%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back of bag for additional precautionary statements.

LEBANON CHEMICAL CORPORATION
LEBANON, PA.



GREENVIEW
EVER PREEN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep out of reach of children and pets. Harmful if swallowed or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact, wash hands with soap and water; flush eyes with running water and get medical attention if irritation persists; wash clothing before reuse.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This two-way product feeds a modern plant food and at the same time prevents broadleaf and grassy weeds from getting started in planting beds. Safe for established evergreens, shrubs and certain listed flowers and vegetables. Do not apply to Ajuga, Myrtle, Pachysandra and other ground covers, as injury may result.

WEEDS CONTROLLED: Annual Bluegrass, Barnyardgrass (Watergrass), Brachiaria, Bromegrass, Cheat, Crabgrass, Foxtail, Goosegrass, Johnsongrass (from seed), Junglerice, Sandbur, Stinkgrass, Sprangletop, Texas Panicum, Wildcane. Also controls these broadleaf weeds: Carelessweed, Carpetweed, Chickweed, Florida Purslane, Goosefoot, Knotweed, Kochia, Lambsquarter, Pigweed, Purslane, Russian Thistle, Stinging Nettle.

EVERGREENS AND SHRUBS

Apply in spring, as evergreen and shrub growth begins, and before weeds sprout. For best results, follow these directions carefully.

1. Before applying, remove all existing broadleaf and grassy weeds, since this product kills sprouting weeds only, not established weeds.
2. Spread Ever Preen evenly. Avoid missing small areas, since weeds will grow in spots not covered. A trowel, scoop or can makes spreading easier.
3. Apply so soil is lightly covered, as shown at right. This represents a rate of 1 lb. per 160 sq. ft.



$\frac{3}{4}$ - 125 sq. ft. registered
.96 lb = 160 sq. ft.

4. For best weed control and proper plant feeding, make a second application 4 weeks after first application. Weed control is increased by lightly raking Ever Preen into soil immediately after application where practical.
5. After application, sprinkle shrubs lightly with water to remove any of the product from foliage.

TREES-ORNAMENTALS-FLOWERS

Ever Preen may be used to control certain weeds in plantings of Roses and other flowers, and around trees and other ornamentals. Your beds will remain cleaner-looking without difficult hand weeding. Remember: Ever Preen does not kill existing weeds; it prevents new weeds from appearing. Ever Preen is not for use with flower seeds.

WHEN TO APPLY

Apply Ever Preen as early as possible in the growing season to take advantage of the season-long weed control it provides. You may apply anytime of the year around shrubs, trees and established plants, and anytime prior to setting transplants.

HOW TO APPLY

Before any application, first remove all existing weeds and lightly cultivate to prepare a smooth soil surface.

For established plants, including flowers at least 2-3 inches tall, sprinkle Ever Preen on the prepared ground at the rate of 1 lb. per 80 sq. ft. Do not apply when foliage is wet. It is advisable to lightly wash into the soil if rain is not imminent.

For pre-planting application, use at the rate of 1 lb. per 80 sq. ft. before setting transplants. Do not use with flower seeds. Gladioli corms less than 1 inch in diameter may be injured by pre-plant application. Do not over-apply. Optimum weed control will result when light watering or rainfall occurs within a few hours of application.

FOR USE ON THESE ESTABLISHED FLOWERS

Achillea, Ageratum, Arctotis, Aster, Balsam, Calendula, California Poppy, Calliopsis, Carnation, Centaurea, Chrysanthemum, Cosmos, Daffodils, Dahlia, Dianthus, Dimorphetheca, Forget-me-not, Four O'Clock, Gaillardia, Gladiolus (corms more than 1" in diameter), Hollyhock, Ixora, (Coccinea), Lobelia, Lupines, Marigold, Morning Glory, Nasturtium, Nicotiana, Periwinkle, Petunia, Phlox, Portulaca, Rudbeckia, Salvia, Scabiosa, Shasta Daisy, Snapdragon, Stocks, Snow on the Mountain, Sunflower, Sweet Alyssum Sweet Pea, Sweet William, Tulips, Zinnia.

VEGETABLES

new 43640-1
2 x
1 lb
per
sq. ft.

Ever Preen may be used for weed control in certain vegetable crops as directed below. Application may be made as much as 10 weeks prior to planting direct seeded vegetables, and anytime prior to setting transplants. Remove all existing weeds first; then lightly cultivate to prepare a smooth soil surface. Rate of application depends on type of soil. Apply at 1 lb. per 320 sq. ft. (18 ft. x 18 ft.) for heavy clay type soils, and at 1 lb. per 640 sq. ft. (25 ft. x 25 ft.) for light soils. Gently rake into the top 1-3 inches of soil within 24 hours of application. Do not use on soils containing more than 10% organic matter.

- (1) From seed: Carrots, Celery, Okra, Southern Peas, Green Peas, Lima Beans, Snap Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Turnip Greens, Mustard Greens, Collard, Kale. Treat prior to seeding.
- (2) Post Emergent: Cantaloupe, Cucumber, Watermelon. Apply after plants are in 3 to 4 true leaf stage.
- (3) Asparagus: Established only. Apply after ferns are removed, but prior to spear emergence.
- (4) Before Transplanting: Celery, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Peppers, Tomatoes. Make application to prepared soil and incorporate into soil prior to placing transplants.
- (5) Potato (except Maine): May be applied after planting, before emergence or after emergence. Use care not to damage seed pieces in cultivating, nor to allow treated soil to contact emerged plant foliage.

In all cases, cultivate the Ever Preen into the soil after application, using care not to damage emerged crops.

LIMITATIONS

Do not apply Ever Preen to muck soils. Do not over-apply; injury may occur to subsequent crops which are sensitive to trifluralin. Do not use with crops other than those listed on this label. Ever Preen is not intended for use on lawns, nor for use in enclosed areas, such as greenhouses.

STORAGE AND DISPOSAL

STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.
PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.
CONTAINER DISPOSAL: Do not reuse container. Discard container in trash.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the face hereof. All statements made concerning this product apply only when used as directed.