


interview in the contraction of the transmission of the trans

the standard and a second plants in the immediate and the

AUTION



## HERBICIDE (2, WITH 12-4-

## FOR PROFE





. L -

Sounday Clu

 $\overline{\mathbf{O}}$ 

ERBIC

D

TIM

N

- 1. Apply uniformally at the rate of 375 lbs. per acre or 8.6 lbs. per 1000 sq. ft. with a fertilizer spreader. This rate gives the recommended herbicide treatment.
- 2. For best results, apply when the weeds are growing actively during the spring and fall seasons. Do not apply to newly seeded areas.
- 3. Increased effectiveness is obtained by applying when the grass is moist such as early in the morning after a dew or following a shower. Avoid, if possible, applying ahead of a heavy rain.
- 4. Do not apply under the dripline of trees or shrubs especially dogwood, willow and locust trees. Avoid drift of dust to susceptible plants. Be careful that water from treated areas does not accumulate around trees or shrubs.
- 5. Do not apply on Dichondra, Centipede, St. Augustine, or Carpet grasses. Temporary injury may be experienced on Bentgrass. Vapors from this product may injure susceptible plants in the immediate vicinity.

#### CAUTION

May cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

If grass clippings from the first mowing after use of this product are removed, do not feed to livestock or use to mulch ornamental shrubs.

Do not contaminate water used for domestic or irrigation purposes.

Do not store near fertilizers, seeds, insecticides or fungicides.

This product is toxic to fish. Keep out of lakes, streams or ponds.

Do not reuse container. Destroy when empty.


SUGGESTED SPREADER SETTINGS					
	Speed		Width	Settings	
Spreader	Ground MPH	PTO RPM	of Coverage	Regular (1#N.)	Half (½#N.)
Jet Spred Cyclone Ezee Flow (109) Lely (HR) Lely (W) Scotts (Drop Type) Scotts (Rotary)	<b>3 3 5 ფ ფ ფ</b>	540 425 425	8 ft. 8 ft. 30 ft. 39 ft. 39 ft. 6 ft.	6 6 A5 711 711 6 <b>M</b>	4½ 4½ C3 5Ⅲ 4½ H

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

#### WARRANTY


The Bishop Company makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. The buyer assumes all risks; including injury or damage, arising out of the storage, handling and/or use of the product when not in strict accordance with directions given herewith.

#### LEBANON CHEMICAL CORPORATION THE BISHOP COMPANY DIVISION LEBANON, PA. 17042



FREICHT BHIPPING BAG Northg registerinth of APPLICABLE PREICHT CLAREFICATION Overeited by UNION CAMP CORPORATION

63548-3 5-61 35593 U-20 POLY LINED



# IERBICIDE (2, WITH 12-4-

1

## WITH 12-4-8 FE

#### 50% OF NITROGEN FROM C

#### MINIMUM GUARANTEED

\*al Nitrogen 'N' 3.6 % Water Insoluble Nitroge

**HN** Country Club ;ES i**5**93 ACCEPTED APP **.** Under the Freiherd Research and a Fungles the training of the first state of the first st