

# CRABGRASS CONTROL

Kills crabgrass  
before it emerges

#### PERCENTAGE OF ACTIVE INGREDIENTS


ACTIVE INGREDIENTS  
Siduron, 1-(2-methylcyclohexyl)-3-phenylurea\* .. 4.00%  
INERT INGREDIENTS..... 95.40%  
EPA REG. NO. 827-223

\*Also known as Tupersin®

2. Rake lawn to remove leaves and other debris.
3. On Established Lawns—apply at the rate of 4 lbs. per 1,000 sq. ft. (10 lb. bag covers 2,500 sq. ft.). Apply after the 1st or 2nd mowing (usually at the time forsythia blossoms start to fall) in the spring before crabgrass and other annual grass weeds emerge.
4. On Newly Seeded Areas—apply at the rate of 2 lbs. per 1,000 sq. ft. (10 lb. bag covers 5,000 sq. ft.). Apply immediately after seeding in early spring before crabgrass and other annual grass weeds emerge.
5. Apply with fertilizer spreader (see settings below) to insure even coverage.
6. If it does not rain for three days after application, water the treated area thoroughly to get desired results. Timing application just ahead of a predicted rain is ideal.

#### CAUTION

May cause irritation of eyes, nose, throat and skin. In case of contact with skin or eyes, flush with plenty of water, for



Agricultural Chemical Division  
Monsanto Company  
St. Louis, Missouri 63103

# before it emerges


## HERBICIDE INGREDIENT STATEMENT

### ACTIVE INGREDIENTS

Siduron, 1-(2-methylcyclohexyl)-3-phenylurea\* .. 4.60%  
 INERT INGREDIENTS..... 95.40%

EPA REG. NO. 827-223

\*Also known as Tuperon®



Agrico Chemical Division  
 Continental Oil Company  
 Memphis, Tennessee 38101

## PREVENT CRABGRASS WHILE AGGREGATING CRABGRASS CONTROL AHEAD TO YOUR LAWN

1. Prevents the emergence of crabgrass seedlings without harming the desirable turfgrasses listed below.
2. Prevents seedlings of Foxtail and Barnyard grass from emerging.
3. Permits safe seeding of your lawn while controlling crabgrass at the same time.

### HOW TO USE

1. For use on newly seeded or established plantings of the following grasses:

Bluegrass	Bentgrass-Strains as follows:
Fescue	Penneross
Redtop	Seaside
Smooth Brome	Highland
Perennial Rye	Astoria
Orchardgrass	Nimisila
Zoysia	C-1, C-7, and C-19

DO NOT USE on other types of grasses or plants listed on this label.

2. Rake lawn to remove leaves and other debris.
3. On Established Lawns—apply at the rate of 4 lbs. per 1,000 sq. ft. (10 lb. bag covers 2,500 sq. ft.). Apply after the 1st or 2nd mowing (usually at the time forsythia blossoms start to fall) in the spring before crabgrass and other annual grass weeds emerge.
4. On Newly Seeded Areas—apply at the rate of 2 lbs. per 1,000 sq. ft. (10 lb. bag covers 5,000 sq. ft.). Apply immediately after seeding in early spring before crabgrass and other annual grass weeds emerge.
5. Apply with fertilizer spreader (see settings below) to insure even coverage.
6. If it does not rain for three days after application, water the treated area thoroughly to get desired results. Timing application just ahead of a predicted rain is ideal.

## CAUTION

May cause irritation of eyes, nose, throat and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

## SUGGESTED SPREADER SETTINGS

Following are recommended spreader settings to get approximately 2 lbs. or 4 lbs. of product per 1,000 sq. ft. of lawn.

SPREADER	SETTING		SPREADER	SETTING	
	2 Lb.	4 Lb.		2 Lb.	4 Lb.
Agrico-Agri-Spred®	5½	7½	Jackson, Seats	5½	7½
Agrico Jet-Spred®			Lawn Beauty	6½	8½
Cyclone	4½	5½	M.T.D.	5½	7½
Central	8	10	Scotts	6	9
Greenfield	8	10			

WALKING SPEED, EVENNESS OF LAWN AND CONDITION OF SPREADER MAY REQUIRE SLIGHTLY DIFFERENT SETTINGS FOR DESIRED COVERAGE.

Satisfaction from the use of this product is guaranteed when used as directed.