

ACCEPTED
 MAR 16 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PESTS REGISTERED
 IN UNDER NO. 271, 229 SUBJECT
 NO ATTACHED COMMENTS.

961-315

Country Club

FERTILIZER

22-10-10 WITH INSECTICIDE

20% OF NITROGEN COMES FROM ORGANIC SOURCES

Use on turf weeds such as: LAWNS
 Annual Blue, Plantain, Cockburn, Shepherd's Purse, Blackwood, Chickweed, Knotweed, Clover, Sheep Sorrel
 and found in golf course fairways, aprons, tees, roughs and on turf (not pasture) areas.

MINIMUM GUARANTEED FERTILIZER ANALYSIS		HERBICIDE INGREDIENT STATEMENT	
Total Nitrogen (N).....	12.0%	Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	0.63%
2.0% Water Insoluble Nitrogen.....		Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	0.32%
Available Phosphate (P ₂ O ₅).....	4.0%	Dimethylamine salt of "Sinox" 2-(2,4,5-trichlorophenoxy) propionic acid.....	0.23%
Soluble Potash (K ₂ O).....	8.0%	Equivalent to 0.15% "Sinox" 2-(2,4,5-trichlorophenoxy) propionic acid.....	
Sulfur (S).....	4.0%	Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....	0.32%
Iron (Fe).....	1.0%	Equivalent to 0.25% 2-(2-methyl-4-chlorophenoxy) propionic acid	
Residual Solids 1100 mg Calcium Carbonate Equivalent per Ton		INERT INGREDIENTS.....	93.82%

EPA Reg. No. E27-229

CAUTION
 KEEP OUT OF REACH OF CHILDREN
 HARMFUL IF SWALLOWED OR INHALED
 SEE BACK OF BAG FOR ADDITIONAL CAUTIONS

T & G PRODUCTS
 DIVISION
RECEIVED
 JAN 25 1972

conoco Agrico Chemical Division
 Continental Oil Company
 Memphis, Tennessee 38101

AgriCo Country Club FERTILIZER

12-0-0 WITH WEED KILLER

HOW TO USE

1. Apply uniformly at the rate of 375 lbs. per acre with a fertilizer spreader. This rate gives the recommended herbicide treatment.
2. For best results, apply when the weeds are growing actively during the spring and fall seasons.
3. Increased effectiveness is obtained by applying when the soil is moist such as early in the morning after a dew or following a shower. When, if possible, applying ahead of a heavy rain.
4. Do not apply under the drip line of trees or shrubs, especially dogwood, willow, and locust trees. Avoid risk of root to root damage plants. Be careful that water from treated areas does not accumulate around the trees.
5. Do not apply on Dichoria, Centipede, St. Augustine, or Carpet grasses. Temporary injury may be expected on Bermudagrass.

CAUTION

May cause irritation of eyes, nose, throat and skin. In case of contact with skin, or eyes, wash with plenty of water; for eyes get medical attention.

If grass clippings from the first mowing after use of this product are removed, do not feed to livestock or use to mulch ornamental shrubs.

Do not store near fertilizers, salts, insecticides, or fungicides.

Do not reuse container. Destroy when empty.

WARRANTY

AgriCo Chemicals, Inc., a subsidiary of The Fertilizer Company, Inc., warrants that this product contains the amount of active ingredients as shown on the label. AgriCo Chemicals, Inc. and its subsidiaries shall not be responsible in any manner for any personal injury, property damage, or loss of crops or other property resulting from the use, storage, or use of this product. The user assumes all risk and liability for any damage or loss resulting from the use of this product on these conditions.