



# GRANULAR CHLORDANE 5

**ACTIVE INGREDIENT:**

Technical Chlordane* .....	5.0%
INERT INGREDIENTS .....	95.0%

\* Equivalent to 3% octachloro-4, 7-methanotetrahydroindane and 2% related compounds.

### DIRECTIONS

**LACCO GRANULAR CHLORDANE 5** is prepared for use against certain pests of crops as stated below. Unless otherwise noted, dosages are in terms of pounds of this product per acre uniformly applied as a broadcast soil treatment.

**CITRUS:** Ants, grasshopper, 100 lbs. as needed as a soil treatment when no fruit is present.

**STRAWBERRY:** Hoplia beetle, brachyrhinus weevil, Fuller's rose beetle, wireworm, 200 lbs. Preplant treatment only. Mix well into upper 4 to 6 inches of soil.

**WIREWORM:** Celery, peppers, potato, sweet potato, 50 to 100 lbs. Preplant treatment only. Mix well into upper 4 to 6 inches of soil.

**ORNAMENTAL LAWN & TURF:** Ant, Asiatic garden beetle, chigger, earwig, European chafer, green June beetle, Japanese beetle, lawn moth, sod webworm, mosquito, scorpion, tick, whitefringed beetle, white grub, wireworm, 50-100 lbs. Water treat area thoroughly after application. Keep children and pets off treated areas until this material has washed into the soil and grass is dry.

## CAUTION!

**KEEP OUT OF REACH OF CHILDREN**

**KEEP CONTAINER CLOSED.**

**DESTROY CONTAINER WHEN EMPTY.**

Do not reuse empty container. • Destroy it by burying with waste or burning. • Stay away from smoke or fumes.

## CAUTION

Harmful if swallowed, inhaled or absorbed through skin or eyes. Do not swallow, inhale or allow contact with skin, eyes or clothing. Wash thoroughly after handling and eating or smoking. Do not contaminate feed and food stuffs. Use only under well ventilated conditions.

This product is toxic to fish, birds, and other wildlife. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important.