

NET WEIGHT: 50 LBS. (XXX Kg.)  
TREATS: XXXX SQ. FT.

# LEBANON COUNTRY CLUB 18-5-9

## FERTILIZER WITH BALAN® FOR TURF

Total Nitrogen (N)	16.00%
5.5% Water Insoluble Nitrogen	
8.5% Ammoniacal Nitrogen	
4.0% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	5.00%
Soluble Potash (K <sub>2</sub> O)	9.00%
Magnesium (Mg)	1.00%
Sulfur (S)	8.00%
Iron (Fe)	1.00%
Manganese (Mn)	0.50%

AC...  
961-321

### ACTIVE INGREDIENT:

\*N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine 1.22%

### INERT INGREDIENTS:

98.78%

TOTAL: 100.00%

\*Balan - a trademark of DowElanco

EPA Reg. No 961-321

(L) (D)  
EPA Est. 961-PA-01.961-IL-01  
Superscript used is first letter of lot number.

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
(See Back Panel for Additional Precautionary Statements)

**MANUFACTURED BY:**  
**Lebanon Chemical Corporation**  
**Lebanon, PA 17042**

Page 1 of 4 Pages  
961-321  
01/16/95

**BEST AVAILABLE COPY**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or spray mist and contact with skin or eyes. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product.

**FIRST AID**

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**IF INHALED:** If discomfort occurs, remove individual to fresh air. Get medical attention if breathing difficulty occurs.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES**

1. This product fertilizes grass and controls smooth and hairy crabgrass, goosegrass, barnyardgrass, green and yellow foxtail and annual bluegrass on established turf of Bluegrass, Ryegrass, Fescuegrass, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass
2. Do not apply in the Spring to bluegrass planted the previous Fall. Do not apply to newly-sprigged areas until root growth is established. Do not apply to bentgrass, dichondragrass, clover lawns or putting greens as possible injury may result.

**BEST AVAILABLE COPY**

3. **Crabgrass, Goosegrass, Barnyardgrass and Foxtail control:**
  - (a) For Warm Season Grasses (Southern), apply this product in early Spring prior to weed grass germination at the rate of 245 lbs. (3 lbs. actual BALAN®) per acre.
  - (b) For Cool Season Grasses, apply in early Spring, prior to weed germination, at the rate of 123 lbs. per acre (1 1/2 lb. BALAN® per acre) for short-season areas, or at 163 lbs. per acre (2 lbs. BALAN® per acre) for longer-season areas.
4. **Annual Bluegrass (Poa Annua) Control:**
  - (a) For Warm Season Grasses, apply this product in early Fall at the rate of 245 lbs. (3 lbs. actual BALAN®) per acre.
  - (b) For Cool Season Grasses, apply this product in early Fall at the rate of 163 lbs. (2 lbs. actual BALAN®) per acre.
5. Reseeding should be delayed approximately 90 days after using Benefin.

### SPREADER SETTINGS

Spreader	Speed		Width of Coverage	Settings		
	Ground MPH	PTO RPM		3 Lbs. Balán®	2 Lbs. Balán®	1 1/2 Lbs Balán®
Jet Spred®	3		8 ft	5	4 1/2	4
Cyclone	3		8 ft	5	4 1/2	4
Ezee Flow (109)	5	540	30 ft.	C3	C2 1/2	C2
Leiy (HR)	5		39 ft.	7II	6II	5 1/2 II
Leiy (W)	5	425	39 ft.	7II	6II	5 1/2 II
Scotts (Drop Type)	3			6	5	4
Scotts (Rotary) RX7	3		10 ft	I	G	F
Scotts (Rotary)	3		8 ft.	J	H	G
Ortho (Drop)	3			6	5	4 1/2
Ortho (Rotary)	3		8 ft.	6	4	3
Lesco (Rotary)	3		10 ft.	J	H	G

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

**BEST AVAILABLE COPY**

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.