

NET WEIGHT 20 LBS.

# Weed Control plus Turf Food with Iron 25-6-6

ACCEPTED  
*April 2, 1973*  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER  
ED UNDER NO. 827-239

*961-324*

### GUARANTEED ANALYSIS\*

TOTAL NITROGEN (N) .....	25.00%	Derived from—Primary Plant Foods: Ammonium Phosphate, Urea, Ureaform, Muriate of Potash.
1.00% Ammoniacal Nitrogen		
8.00% Water Insoluble Nitrogen		
15.40% Water Soluble Nitrogen		
AVAILABLE PHOSPHORIC ACID (P <sub>2</sub> O <sub>5</sub> ) .....	6.00%	Derived from—Micro Nutrient Plant Foods: Chelates, Frits, Sulfates and Oxides.
**SOLUBLE POTASH (K <sub>2</sub> O) .....	6.00%	
COPPER (Cu) .....	.10%	
IRON (Fe) .....	1.00%	*In Canada and Michigan, the heading reads "MINIMUM GUARANTEED ANALYSIS".
MANGANESE (Mn) .....	.27%	
ZINC (Zn) .....	.20%	**In states requiring, reads "WATER SOLUBLE (K <sub>2</sub> O)".
Potential Acidity:		
900 lbs. Calcium Carbonate Equivalent per ton.		

ACTIVE INGREDIENTS .....	1.47%
Isocetyl ester of Silvex	
[2-(2, 4, 5-Trichlorophenoxy) Propionic acid]*** .....	0.47%
Dimethylamine salt of 2, 4-Dichlorophenoxyacetic acid**** .....	1.00%
INERT INGREDIENTS .....	98.53%
EPA REG. NO. 827-239- <del>XXXX</del>	Total 100.00%

\*\*\*Equivalent to 0.31% [2-(2, 4, 5-Trichlorophenoxy) Propionic acid], or 0.54 lbs. per acre.  
\*\*\*\*Equivalent to 0.83% 2, 4-Dichlorophenoxyacetic acid (2, 4-D), or 1.44 lbs. per acre.

AGRICO CHEMICAL COMPANY  
TURF AND GARDEN DIVISION  
BOX 3493  
TULSA, OK. 74101

## DIRECTIONS FOR USE

### FOR BEST RESULTS:

Apply when grass is wet with dew, or wet from sprinkling. This will help the weedkiller chemicals stick to the weed surfaces where they do their work. DO NOT WATER FOR 3 DAYS AFTER APPLICATION—this would only wash the weedkiller chemicals off the weed surface. Do not mow for several days before and after application.

WHEN TO USE: In spring, late summer or fall, when weeds are growing actively and temperature is at or above 60°F.

### WEEDS CONTROLLED:

Broadleaf weeds controlled	Lawn pennywort	White clover
Dandelion	Pigweed	Henbit
Ragweed	Ground Ivy	Thyme-leaved speedwell
Plantain	Sow Thistle	Yarrow
Buckhorn	Vining-type weeds controlled	Knotweed
Burdock	Common chickweed	Spotted spurge
Mustard	Mouse ear chickweed	Black medic
Nettle	Little hop clover	Purslane
Jimson weed	Suplover	Lippia

RATE: This bag will treat 5,000 sq. ft. (for smaller areas, use 4 lbs. per acre)

## DIRECTIONS FOR USE

### FOR BEST RESULTS:

Apply when grass is wet with dew, or wet from sprinkling. This will help the weedkiller chemicals stick to the weed surfaces where they do their work. **DO NOT WATER FOR 3 DAYS AFTER APPLICATION**—this would only wash the weedkiller chemicals off the weed surface. Do not mow for several days before and after application.

**WHEN TO USE:** In spring, late summer or fall, when weeds are growing actively and temperature is above 60°F.

### WEEDS CONTROLLED:

<u>Broadleaf weeds controlled</u>	<u>Lawn pennywort</u>	<u>White clover</u>
<u>Dandelion</u>	<u>Pigweed</u>	<u>Hanbit</u>
<u>Ragweed</u>	<u>Ground Ivy</u>	<u>Thyme-leaved speedwell</u>
<u>Plantain</u>	<u>Sow Thistle</u>	<u>Yarrow</u>
<u>Buckhorn</u>	<u>Vining-type weeds controlled</u>	<u>Knotweed</u>
<u>Burdock</u>	<u>Common chickweed</u>	<u>Spotted spurge</u>
<u>Mustard</u>	<u>Mouse ear chickweed</u>	<u>Black medic</u>
<u>Nettle</u>	<u>Little hop clover</u>	<u>Purslane</u>
<u>Jimson weed</u>	<u>Bur-clover</u>	<u>Lippia</u>

**RATE:** This bag will treat 5,000 sq. ft. (for smaller areas, use 4 lbs. per 1,000 sq. ft. of lawn area.) See spreader setting chart. **DO NOT APPLY MORE THAN RECOMMENDED RATE — EXCESS APPLICATION WILL NOT IMPROVE RESULTS.**

**IMPORTANT:** Use only on established lawns. Avoid drift of dust on vegetables, flowers, ornamental trees and shrubs, and other desirable plants. If some of the material comes in contact with desirable plants, it is recommended that the plants be syringed thoroughly with water as soon as possible. Vapors from this product may injure susceptible plants in the immediate vicinity.

**DO NOT STORE** near other fertilizer, seed, insecticides or fungicides.

**DO NOT USE** on St. Augustine grass, carpetgrass, centipede, bentgrass, and Dichondra as possible injury may result.

SPREADER SETTINGS (1 lb. per 250 sq. ft. rate)	
PENNCRAFT	5½
CENTRAL QUALITY	5½
CYCLONE	4
LAWN BEAUTY-MODERN	5
JACKSON	5
SCOTT	5½
These settings are approximate. Settings vary with model, age and condition of spreader.	

**CAUTION:** Avoid inhalation of dust or contact with skin, eyes or clothing. Wash thoroughly after handling. If swallowed induce vomiting and consult your physician.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply in any manner not specified on the label. Do not use where runoff will contaminate water used for irrigation or domestic use.

Do not reuse container. Destroy by burying in a safe place away from water supplies. Do not burn.

816-0108