

NET WEIGHT: 20 LBS. (9.07 Kg)
COVERS: 5,000 OR 10,000 SQ. FT.

GREEN GOLD

INSECT AND GRUB CONTROL

WITH DURSBAN*

5,000 Sq. Ft. Coverage: Kills Soil Feeding Insects - Japanese Beetle Larvae, European
Chafer Larvae
10,000 Sq. Ft. Coverage: Kills Soil Surface Insects - Sod Webworms, Cutworms, Chinch
Bugs, Ants, Mole Crickets, Brown Dog
Ticks, Earwigs, Fleas, Chiggers, Clover
Mites, Grasshoppers, Armyworms

ACTIVE INGREDIENT:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate]* 1.14%

INERT INGREDIENTS: 98.86%

TOTAL: 100.00%

*Dursban - a trademark of DowElanco

EPA Reg. No. 961-326

(L) (D)
EPA Est. 961-PA-01.961-IL-01
Superscript used is first letter of lot number.

CAUTION
KEEP OUT OF REACH OF CHILDREN
See Back Panel for Additional Precautionary Statements and Statement of
Practical Treatment

ACCEPTED
APR 18 1995
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 961-326

MANUFACTURED BY
Lebanon Chemical Corporation
Lebanon, PA 17042

Page 1 of 4 Pages
961-326
01/16/95

1084

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

May be harmful if swallowed.

Do not take internally. Avoid inhaling dust. Avoid eye contact. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air if symptoms of cholinesterase inhibition appear, and get medical attention immediately.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection, is an antidote.

ENVIRONMENTAL HAZARD

This product is toxic to fish, birds and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

5,000 SQ. FT. FOR SOIL GRUBS

For the control of larvae of Japanese Beetle and European Chafer, apply at the rate of 4 to 8 lbs. per 1,000 sq. ft. Immediately wash into the soil with 1/2 to 1 inch of water. Treat when white grubs are young and actively feeding near the soil surface, usually during late July and August, or as recommended by your local Agricultural Extension Service Specialist.

10,000 SQ. FT. FOR SURFACE INSECTS

For Sod Webworms, Cutworms, Chinch Bugs, Ants, Mole Crickets, Brown Dog Ticks, Fleas, Chiggers, Clover Mites, Grasshoppers, Armyworms and Earwigs, apply this product at the rate of 2 lbs. per 1,000 sq. ft. Treat when insects first appear. Immediately after application, sprinkle the turf with water to wash granules into the zone of insect infestation. Repeat as necessary.

Keep children and pets off treated areas until the material is washed into the soil and grass is dry.

Do not mow for 24 hours following treatment.

Use a fertilizer spreader to insure an even application. Follow the suggested spreader settings.

SPREADER SETTINGS

2 Lb. Setting

Show 'N Tell	7	Glorion	7
Greenfield	7	Queen Turf	7
Scott	4	Sears	7
Central	7	Jet Spred	3 1/2
M.T.D.	2 1/2	Cyclone	3 1/2
Jackson	5 1/2	Scott Rotary	D

4 Lb. Setting

Show 'N Tell	8	Glorion	8
Greenfield	8	Queen Turf	8
Scott	8 1/2	Sears	8
Central	8	Jet Spred	4 1/2
M.T.D.	3 1/2	Cyclone	4 1/2
Jackson	6 1/2	Scott Rotary	K

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 3 1/2 oz. at 2 lb. setting, 6 1/2 oz. at 4 lb. setting. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

STORAGE AND DISPOSAL

STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.