



6

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.

101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200.

PTEN 9/11

8/11

PTEN 9/11

PTEN 9/11

PTEN 9/11

NET WEIGHT 20 LBS.

Weed Control

plus

Turf Food

with iron

25-6-6

GUARANTEED ANALYSIS*

| | |
|---|---|
| <p>TOTAL NITROGEN (N)25.00%</p> <p> 1.00% Ammoniacal Nitrogen</p> <p> 8.00% Water Insoluble Nitrogen</p> <p> 15.40% Water Soluble Nitrogen</p> <p>AVAILABLE PHOSPHORIC ACID (P₂O₅) 6.00%</p> <p>◆◆ SOLUBLE POTASH (K₂O) 6.00%</p> <p>COPPER (Cu)10%</p> <p>IRON (Fe) 1.00%</p> <p>MANGANESE (Mn)27%</p> <p>ZINC (Zn)20%</p> <p>CHLORINE (Cl) (Maximum) 5.00%</p> <p>Potential Acidity: 980 lbs. Calcium Carbonate Equivalent per ton.</p> | <p>Derived from—Primary Plant Foods: Ammonium Phosphate, Urea, Ureaform, Muriate of Potash.</p> <p>Derived from—Micro Nutrient Plant Foods: Chelates, Frits, Sulfates and Oxides.</p> <p>*In Canada and Michigan, the heading reads "MINIMUM GUARANTEED ANALYSIS".</p> <p>◆◆ In states requiring, reads "WATER SOLUBLE (K₂O)".</p> |
|---|---|

| | |
|---|---------------|
| ACTIVE INGREDIENTS..... | 1.20% |
| Isocetyl ester of Silvex | |
| [2-(2, 4, 5-Trichlorophenoxy) Propionic acid]*** | 0.44% |
| Dimethylamine salt of 2, 4-Dichlorophenoxyacetic acid**** | 0.76% |
| INERT INGREDIENTS..... | 98.80% |
| EPA Reg. No. 827-241 8000 | Total 100.00% |

***Equivalent to 0.31% [2-(2, 4, 5-Trichlorophenoxy) Propionic acid], or 0.54 lbs. per acre.
 ****Equivalent to 0.63% 2, 4-Dichlorophenoxyacetic acid (2, 4-D), or 1.09 lbs. per acre.

AGRICO CHEMICAL COMPANY
 TURF AND GARDEN DIVISION
 BOX 3493
 TULSA, OK. 74101

589-1010

DIRECTIONS FOR USE

FOR BEST RESULTS:

Apply when grass is wet with dew, or wet from sprinkling. This will help the weedkiller chemicals stick to the weed surfaces where they do their work. DO NOT WATER FOR 3 DAYS AFTER APPLICATION—this would only wash the weedkiller chemicals off the weed surface. Do not mow for several days before and after application.

WHEN TO USE: In spring, late summer or fall, when weeds are growing actively and temperature is above 60°F.

WEEDS CONTROLLED:

| | | |
|---|--|---|
| <p><u>Broadleaf weeds controlled</u></p> <p><u>Dandelion</u></p> <p><u>Ragweed</u></p> <p><u>Plantain</u></p> <p><u>Buckhorn</u></p> <p><u>Burdock</u></p> <p><u>Mustard</u></p> <p><u>Nettle</u></p> <p><u>Jimson weed</u></p> | <p><u>Lawn pennywort</u></p> <p><u>Pigweed</u></p> <p><u>Ground Ivy</u></p> <p><u>Sow Thistle</u></p> <p><u>Vining-type weeds controlled</u></p> <p><u>Common chickweed</u></p> <p><u>Mouse ear chickweed</u></p> <p><u>Little hop clover</u></p> <p><u>Bur clover</u></p> | <p><u>White clover</u></p> <p><u>Henbit</u></p> <p><u>Thyme-leaved speedwell</u></p> <p><u>Yarrow</u></p> <p><u>Knotweed</u></p> <p><u>Spotted spurge</u></p> <p><u>Black medic</u></p> <p><u>Purslane</u></p> <p><u>Lippia</u></p> |
|---|--|---|

RATE: This bag will treat 5,000 sq. ft. (for smaller areas, use 4 lbs. per 1,000 sq. ft. of lawn area.) See spreader setting chart. DO NOT APPLY

EPA Reg. No. 627-241-0000
 ***Equivalent to 0.51% [2-(2, 4, 6-Trichlorophenoxy) Propionic acid], or 0.84 lbs. per acre.
 ***Equivalent to 0.63% 2, 4-Dichlorophenoxyacetic acid (2, 4-D), or 1.09 lbs. per acre.

AGRICO CHEMICAL COMPANY
 TURF AND GARDEN DIVISION
 BOX 3493
 TULSA, OK. 74101

589-1010

DIRECTIONS FOR USE

FOR BEST RESULTS:

Apply when grass is wet with dew, or wet from sprinkling. This will help the weedkiller chemicals stick to the weed surfaces where they do their work. DO NOT WATER FOR 3 DAYS AFTER APPLICATION—this would only wash the weedkiller chemicals off the weed surface. Do not mow for several days before and after application.

WHEN TO USE: In spring, late summer or fall, when weeds are growing actively and temperature is above 60°F.

WEEDS CONTROLLED:

Broadleaf weeds controlled

Dandelion
Ragweed
Plantain
Buckhorn
Burdock
Mustard
Nettle
Jimson weed

Lawn pennywort

Ploweed
Ground ivy
Sow Thistle
Vining-type weeds controlled
Common chickweed
Mouse ear chickweed
Little hop clover
Bur clover

White clover

Henbit
Thyme-leaved speedwell
Yarrow
Knotweed
Spotted spurge
Black medic
Purslane
Lippie

RATE: This bag will treat 5,000 sq. ft. (for smaller areas, use 4 lbs. per 1,000 sq. ft. of lawn area.) See spreader setting chart. DO NOT APPLY MORE THAN RECOMMENDED RATE — EXCESS APPLICATION WILL NOT IMPROVE RESULTS.

IMPORTANT: Use only on established lawns. Avoid drift of dust on vegetables, flowers, ornamental trees and shrubs, and other desirable plants. If some of the material comes in contact with desirable plants, it is recommended that the plants be syringed thoroughly with water as soon as possible. Vapors from this product may injure susceptible plants in the immediate vicinity.

DO NOT STORE near other fertilizer, seed, insecticides or fungicides.

DO NOT USE on St. Augustine grass, carpetgrass, centipede, bentgrass, and Dichondra as possible injury may result.

| SPREADER SETTINGS (1 lb. per 250 sq. ft. rate) | |
|--|----|
| PENNCRAFT | 5½ |
| CENTRAL QUALITY | 5½ |
| CYCLONE | 4 |
| LAWN BEAUTY-MODERN | 5 |
| JACKSON | 5 |
| SCOTT | 5½ |
| These settings are approximate. Settings vary with model, age and condition of spreader. | |

CAUTION: Avoid inhalation of dust or contact with skin, eyes or clothing. Wash thoroughly after handling. If swallowed, induce vomiting and consult your physician. Avoid contamination of foodstuffs.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply in any manner not specified on the label. Do not use where runoff will contaminate water used for irrigational or domestic use.

Do not reuse container. Destroy by burying in a safe place away from water supplies. Do not burn.

816-0136