

LEBANON GREEN GOLD® GARDEN WEED PREVENTER

CONTAINS TRIFLURALIN

FOR HOME USE ONLY

NOT FOR USE ON LAWNS

Prevents appearance of the listed weeds in home flower and home vegetable garden beds by killing their seeds as they germinate.

ACTIVE INGREDIENT:

Trifluralin: a,a,a-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	1.75%
OTHER INERT INGREDIENTS:	<u>98.25%</u>
TOTAL	100.00%

EPA Reg. No. 961-335

(L) (D)
EPA Est. 961-PA-01:961-IL-01
Superscript used is first letter of lot number.


**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Additional Precautionary Statements.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.lebsea.com

NET WEIGHT: 13 LBS. 5 OZ. (6.05 kg)

**COVERAGE: 10,000 SQ. FT. for Heavy Soil
20,000 SQ. FT. for Light Soil**



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing particulate material and contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove clothing and wash before reuse. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

FIRST AID

IF IN EYES: Flush eyes with plenty of water. Call a physician.

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation develops.

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF INHALED: Remove individual to fresh air. If breathing difficulty occurs, get medical attention. If not breathing, provide cardiopulmonary resuscitation (mouth-to-mouth) and get medical attention immediately.

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner which will expose canals, lakes, streams, ponds, marshes, or estuaries to aerial drift, directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

Do not apply directly to water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.

PESTICIDE DISPOSAL: Securely wrap unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

GENERAL INFORMATION

HOW TO APPLY

On garden areas being prepared for subsequent planting, first loosen soil thoroughly, remove all visible weeds and smooth surface. Apply this product at the recommended rate over the surface uniformly, and rake or spade the product into the soil to a depth of 2 to 3 inches. Subsurface incorporation should be done in two directions, at right angles to each other, to insure uniform distribution of particles. Shallow cultivation during the growing season will not reduce effectiveness.

On established plantings, remove all visible weeds, and then apply this product at the recommended rate uniformly over the soil surface. Rake the product into the soil, being careful not to disturb the established plants or roots.

Allow untreated area for planting of corn, squash and other vegetables not specified for treatment.

WEEDS CONTROLLED

Annual Grasses

Crabgrasses
Barnyardgrass (Watergrass)
Foxtails (including Giant Foxtail)

Johnsongrass (from seed)
Goosegrass
Wild Cane (Shattercane)

Texas panicum
Stinkgrass
Bromegrass

Brachiaria
Sandbur
Jungle-rice

Sprangletop
Cheat

3 9 3

Annual Broadleaves

Pigweeds (*Spiny, Redroot*)
Carelessweed

Lambsquarters
Carpetweed

Russian Thistle
Kochia

Purslane
Florida Pusley

Knotweed
Stinging Nettle

Goosefoot
Chickweed

WILL NOT CONTROL certain resistant weeds such as Cocklebur, Velvetleaf, Jimsonweed, Ragweed, Venice mallow, Nutgrass and established Bermudagrass and Johnsongrass.

APPLICATION RATES

One pound of this product is equal to 2 4/5 cups.

Use the following chart to aid in the proper application at the various recommended rates:

Per 100 Sq. Ft.	1 Lb./1500 Sq. Ft. 3 Tablespoons	1 Lb./1000 Sq. Ft. 4 1/2 Tablespoons	1 Lb./750 Sq. Ft. 6 Tablespoons
--------------------	-------------------------------------	---	------------------------------------

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES. FOR USE ON PLANTS INTENDED FOR AESTHETIC PURPOSES OR CLIMATIC MODIFICATIONS AND BEING GROWN IN INTERIOR PLANTSCAPES, ORNAMENTAL GARDENS OR PARKS OR ON GOLF COURSES OR LAWNS AND GROUNDS.

ENTRY RESTRICTIONS

Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

This product is not recommended for use on muck soils. Do not apply to any vegetable crop not listed on the label. Do not apply to Ajuga, Myrtle, Pachysandra and other ornamental ground covers as injury may result. Use only at the rate recommended; overtreatment may cause injury. Applications made according to directions under normal growing conditions will not harm treated plants.

Wet weather, including disease, cold weather in combinations with deep planting and excessive moisture, high salt concentration and drought may weaken crop seedlings and increase the possibility of damage from this product. Under these conditions, delayed crop development may result.

APPLICATION INSTRUCTIONS

FOR VEGETABLES

RECOMMENDED FOR USE ON Drybeans, Southern peas (*cowpeas, field peas, black-eyed peas*), Snap-beans, Lima beans, Okra and transplants of Tomatoes, Peppers, Broccoli, Brussels Sprouts, Cabbage, Cauliflower. Apply from 6 weeks before planting up to day of planting. Apply at the following rates, **except on Snap beans and Lima beans:** Use 1 lb. to 1,500 sq. ft. on sandy and sandy loam (light) soils. Use 1 lb. to 1,000 sq. ft. on loam (medium) soils. Use 1 lb. to 750 sq. ft. on silt and clay (heavy) soils. On sandy soils containing 2%-5% organic matter, apply 1 lb. per 1,000 sq. ft. and on sandy soils containing 5%-10% organic matter, apply 1 lb. per 750 sq. ft.

On Snap beans and Lima beans, apply 1 lb. to 1,500 sq. ft. on sandy, sandy loam or loam soils and 1 lb. to 1,000 sq. ft. on silt and clay soils.

FOR ORNAMENTAL SHRUBS, TREES AND FLOWERS

RECOMMENDED FOR USE ON Established Ornamental Shrubs and Trees and the Following Flowers: Ageratum, Aster, Balsam, Calendula, California Poppy, Calliopsis, Carnation, Cosmos, Daffodils, Dahlia, Forget-Me-Not, Four-O'Clocks, Gaillardia, Gladiolus, Hollyhock, Lobelia, Marigold, Morning Glory, Nasturtium, Periwinkle, Petunia, Portulaca, Shasta Daisy, Snapdragon, Stocks, Sunflower, Sweet Alyssum, Sweet Pea, Tulips, Zinnia. **Apply at the rate of 1 lb. to 750 to 1,000 sq. ft. in the Spring around shrubbery and trees and in flower beds after plants have become established.**

Hand held granular applicators designed to apply herbicide or insecticide granules are ideal for applying this product. Calibrate the applicator according to the manufacturer's directions prior to use and check frequently during application to be sure the equipment is working properly and distributing the granules uniformly. Do not apply more than the recommended amount.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

