


(FRONT)

Net Wt. 15 lb.

Covers 5,000 sq. ft.

LEBANON

Weed Control with Fertilizer
32-3-3



This product is a remarkably effective lawn fertilizer which combines fast and long-lasting feeding action, plus effective weed control. This product produces a thick, green turf while killing weeds to make way for new turf growth.

Lebanon Weed Control with Fertilizer 32-3-3 is designed for the control of dandelion, clover, and many other broadleaf weeds. Weeds contacted by this product gradually die and disappear from the lawn. See back panel for a list of weeds controlled.

| ACTIVE INGREDIENTS | BY WEIGHT |
|--|---------------|
| *Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid | 1.35% ✓ |
| Isooctyl Ester of 2-(2-methyl-4-chlorophenoxy)Propionic Acid** | 1.37% ✓ |
| INERT INGREDIENTS | <u>97.28%</u> |
| TOTAL | 100.00% |

*2,4-Dichlorophenoxyacetic Acid Equivalent .90% Isomer specific by AOAC Method.

**Equivalent to .90% 2-(2-methyl-4-chlorophenoxy)Propionic Acid.

EPA Reg. No. 961-()

Keep Out of Reach of Children.

CAUTION

See back panel for additional precautionary statements.

LEBANON CHEMICAL CORPORATION
Lebanon, Pa. 17042

(BACK-1)

Net Weight 15 lbs.

Covers 5,000 sq. ft.

Lebanon
Weed Control with Fertilizer
32-3-3

Guaranteed Fertilizer Analysis

| | |
|-----------------------------------|-------|
| Total Nitrogen (N) | 32.0% |
| 2.0% Ammoniacal Nitrogen | |
| 6.0% Water Insoluble Nitrogen | |
| 25.0% Water Soluble Urea Nitrogen | |
| Available Phosphoric Acid (P2O5) | 3.0% |
| Soluble Potash (K2O) | 3.0% |

Primary nutrients derived from ammonium phosphate, urea, methylene ureas, muriate of potash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes, or clothing.
Avoid breathing dust.
Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is designed for control of broadleaf weeds in lawns such as: BINDWEED, BLACK MEDIC, BLUE LETTUCE, BUCKHORN, BURDOCK, CHICKWEED, CHIRY, CINQUEFOIL, CLOVER, COCKLEBUR, CROTON, DANDELION, DOCK, FANWEED, GALINSOGA, GOLDENROD, GROUND IVY, HAWKWEED, HEAL-ALL, HENBIT, JEWELWEED, JIMSON WEED, KOCHIA, KNOTWEED, LAMBSQUARTERS, MALLOW, MORNING GLORY, MUSTARD, NETTLES, OXALIS, PENNYWORT, PEPPER GRASS, PIGWEED, PLANTAIN, POVERTY WEED, PURSLANE, RAGWEED, SHEPHERD'S PURSE, SMARTWEED, SPEEDWELL, SPOTTED SPURGE, THISTLE,

(BACK-2)

VERVAINS, VETCH, WILD ASTER, WILD LETTUCE, WILD RADISH, and YARROW.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THE FOLLOWING CAREFULLY BEFORE APPLICATION

This product may be used on Blue Grass, Fescue, Rye, Bermuda, Bahia, and Zoysia lawns.

DO NOT USE on Dichondra, Carpet grass and St. Augustine grasses.

DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables or ornamental shrubbery.

WHEN TO APPLY: Best results are obtained when weeds are actively growing in spring or early fall, at temperatures of 60° F or higher, but may be effectively used anytime during the growing season.

IMPORTANT: Apply when grass is wet from recent rain, sprinkling or dew. Do not apply to newly seeded areas until grass has been mowed two or three times. Do not water treated area or apply within two days of expected rainfall. Do not reseed area for at least four weeks after application. Do not over or under apply - use at recommended rate. Keep children and pets off treated area for several days, until product has been washed into soil by subsequent rainfall or sprinkling.

Avoid contact with valuable shrubs, plants or flowers. Vapors from this product may injure susceptible plants in the immediate vicinity.

HOW TO APPLY: Apply at the rate of 3 lbs. per 1,000 sq. ft. (15 lb. bag covers 5,000 sq. ft.)

Lawns that are heavily overrun by weeds and that are in poor condition may require more than one application of this product. A suggested second application should be made in approximately 30 days. Lawn can be reseeded 4 to 5 weeks after weeds are killed.

Apply evenly with a lawn spreader to assure uniform distribution and correct coverage of area to be treated.

SUGGESTED SPREADER SETTINGS

NOTE: These settings are approximate. Check your spreader on 100 square feet and adjust to apply 4 3/4 ounces per 100 sq. ft. Rate of walk, moisture and

(BACK-3)

spreader condition can vary the setting necessary to apply the correct amount. When using a spinner type (rotary) spreader, use extreme caution to avoid any contact with desirable plants.

| | | | | | |
|-------------|---|------------|----|--------------|---|
| Show-N-Tell | 6 | Sears | 5 | Cyclone | 3 |
| Central | 6 | Scott drop | 4 | Jet Spread | 3 |
| Jackson | 6 | Glorion | 5½ | Scott Rotary | E |

STORAGE

Do not store near food, feed, seed, or other pesticides.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

WARRANTY

Lebanon Chemical Corporation warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

LEBANON CHEMICAL CORPORATION
Lebanon, Pa. 17042